

Fermion masses and mixing in models with $SO(10) \times A_4$ symmetry

Federica Bazzocchi¹, Michele Frigerio² and Stefano Morisi¹

¹ *AHEP Group, Instituto de Física Corpuscular – C.S.I.C./Universitat de València
Edificio Institutos de Paterna, Apt 22085, E-46071 Valencia, Spain*

² *Institut de Physique Théorique, CEA-Saclay, 91191 Gif-sur-Yvette Cedex, France*

We study the flavour sector in models where the three families of matter are unified in a $(16, 3)$ representation of the $SO(10) \times A_4$ group. The necessary ingredients to realize tri-bi-maximal mixing in the lepton sector are identified systematically. The non-renormalizable operators contributing to the fermion mass matrices play an important role. We also present a mechanism to explain the inter-family mass hierarchy of quarks and charged leptons, which relies on a ‘universal seesaw’ mechanism and is compatible with tri-bi-maximal mixing.

I. INTRODUCTION

Before the experimental determination of neutrino oscillation parameters, a common theoretical prejudice was the expectation of small lepton mixing angles, by analogy with the CKM angles in the quark sector. This was thought to be a generic prediction of Grand Unification Theories, as they incorporate some form of quark-lepton symmetry. On the contrary, experiments have shown that the 2 – 3 lepton mixing is close to maximal, the 1 – 2 mixing is large, while the 1 – 3 angle can be as large as the Cabibbo angle [1, 2]. With the benefit of hindsight, this disparity between quarks and leptons was interpreted as a manifestation of the Majorana nature of neutrinos. From then on, many grand unified models have been built, in which large lepton mixing angles appear naturally (see e.g. [3, 4, 5, 6, 7]).

Several conjectures on special ‘symmetric’ values of the lepton mixing angles have been also proposed. For example, the bi-maximal mixing pattern ($\theta_{23} = \theta_{12} = \pi/4$ and $\theta_{13} = 0$) was extensively studied but it is by now excluded, since solar neutrino experiments require $\theta_{12} = (35 \pm 4)^\circ$ at $3\text{-}\sigma$. Current data are in very good agreement, instead, with the so-called tri-bi-maximal (TBM) mixing scheme [8], where the second and third neutrino mass eigenstates have the following flavour content: $\nu_2 = (\nu_e + \nu_\mu + \nu_\tau)/\sqrt{3}$ and $\nu_3 = (\nu_\mu - \nu_\tau)/\sqrt{2}$. Such special values of the angles may point to a non-abelian family symmetry. In particular, the TBM scheme seems to require that the three families of lepton doublets belong to the same dimension three representation, the simplest suitable symmetry being the discrete group A_4 [9, 10] (early applications of A_4 as a family symmetry are discussed in [11, 12]).

While generically large lepton mixing angles found their place in the framework of Grand Unification, it is a much tougher task to justify special values of such angles. The first few attempts to build unified models with TBM mixing have encountered several difficulties and concrete realizations so far require many degrees of complication. In most A_4 models for TBM mixing, the lepton doublets and singlets transform differently under the family symmetry, so that the allowed embedding is $SU(5)$ unification [13, 14, 15], or the Pati-Salam group [16]. The alternative option, to take all leptons transforming as triplets under A_4 , was later considered [17] and it was shown [18] to be suitable to achieve TBM mixing. This assignment is compatible with $SO(10)$ unification of the gauge interactions. Few models based on the $SO(10) \times A_4$ group

have been studied [19, 20]. A different scheme to achieve TBM mixing in $SO(10)$ unification makes use of an $SU(3)$ family symmetry [21] or its discrete subgroup $\Delta(27)$ [22].

In this paper, we investigate systematically the structure of fermion masses and mixing in models with $SO(10)$ unification and A_4 family symmetry. This is the minimal framework for a complete unification of the three families of matter, since the three $SO(10)$ spinors of dimension 16 can transform as a triplet under A_4 . The scheme for realizing TBM lepton mixing consists of breaking A_4 into its Z_3 and Z_2 subgroups in the charged lepton and neutrino sectors, respectively. This misalignment results in TBM values for the lepton mixing angles. On the other hand, the flavour alignment of the up and down quark sectors should be approximately maintained, in order to explain the smallness of the CKM angles. Since in our framework all matter fields belong to the same representation of $SO(10) \times A_4$, the Yukawa couplings of the different sectors are strictly related and it is particularly challenging to satisfy all constraints.

More specifically, one faces the need to generate two independent structures for the up-quark mass matrix M_u and the neutrino Dirac mass matrix M_ν , which in $SO(10)$ are strictly related. It has been shown [19] that a sufficiently complicated arrangement of higher dimensional operators can disentangle the two structures. Another way out [20] is to assume that the light neutrino mass matrix, m_ν , is independent from the form of M_ν and it is instead generated by the coupling of two lepton doublets to a Higgs triplet. However, in the context of $SO(10)$ theories this assumption cannot be realized exactly, as discussed later. Another difficulty of A_4 models, which becomes even more severe in $SO(10)$, is the need to introduce non-vanishing CKM parameters without generating too large deviations from the TBM lepton mixing. We will reconsider these issues, generalizing previous results and identifying new solutions.

Perhaps the most serious shortcoming of A_4 flavour models is that the hierarchy between the masses of the three families can be accommodated, but is not explained. This problem can be addressed [23] in A_4 models where lepton doublets transform as $L_i \sim 3$ while charged lepton singlets transform as $e_i^c \sim 1, 1', 1''$. In this case one can introduce an extra family symmetry that distinguishes the three families. For example, a $U(1)$ family symmetry of the Froggatt-Nielsen type [24], with different charges for the three e_i^c , can explain the hierarchy of the charged lepton masses. This mechanism can be promptly extended in an $SU(5)$ -invariant fashion to the down and up quarks. The origin of the inter-family hierarchy is more problematic in $SO(10)$, since both chiralities of matter transform as A_4 triplets. As a consequence, a universal mass term of the type $m\psi_i\psi_i^c$ is allowed [25] and no extra symmetry can distinguish the families. We will argue that a natural solution for this problem emerges from the mixing with heavy vector-like families. The hierarchy arises in the quark and charged lepton sectors precisely because there A_4 is broken to Z_3 and each family transforms differently under this residual symmetry.

The paper is organized as follows. In section II (and in appendix A) we recall the structure of the fermion mass matrices which lead to TBM mixing in A_4 models. In section III (and in appendix B 1) we study $SO(10) \times A_4$ models with renormalizable Yukawa couplings and investigate the possibility to realize the TBM mixing scheme. In section IV (and in appendix B 2) we extend our analysis to dimension five operators, which overcomes several difficulties of the renormalizable models. In section V we present the mechanism to explain the mass hierarchy between the three families, in a way that is compatible with TBM mixing. In section VI we summarize our main results.

II. THE ROUTE TO TRI-BI-MAXIMAL MIXING

In this section we review a scheme to achieve TBM lepton mixing, that was employed in the construction of several A_4 models [23, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39] and turns out to be the one

suitable in the context of $SO(10)$ unification [18, 19, 20, 40].

Let us introduce a set of simple mass matrix structures which can accommodate all fermion masses, leading at the same time to TBM lepton mixing and vanishing quark mixing. We adopt a supersymmetric notation, with $-\mathcal{L}_m = uM_u u^c + dM_d d^c + eM_e e^c + \frac{1}{2}\nu m_\nu \nu$, where

$$M_f = \begin{pmatrix} A_f & B_f & C_f \\ C_f & A_f & B_f \\ B_f & C_f & A_f \end{pmatrix}, \quad f = u, d, e, \quad m_\nu = \begin{pmatrix} a & 0 & b \\ 0 & c & 0 \\ b & 0 & a \end{pmatrix}. \quad (1)$$

These charged fermion and neutrino Majorana mass matrices can be diagonalized as follows:

$$\begin{aligned} D_f &= U_f^T M_f U_f^* = \text{diag}(A_f + B_f + C_f, A_f + \omega^2 B_f + \omega C_f, A_f + \omega B_f + \omega^2 C_f) \equiv \text{diag}(m_{f1}, m_{f2}, m_{f3}), \\ d_\nu &= U_\nu^T m_\nu U_\nu = \text{diag}(a + b, c, a - b) \equiv \text{diag}(m_1, m_2, m_3), \end{aligned} \quad (2)$$

where $\omega \equiv e^{2\pi i/3} = (-1 + \sqrt{3}i)/2$ and the unitary mixing matrices are given by

$$U_f = U_\omega \equiv \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}, \quad U_\nu = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & -1 \\ 0 & \sqrt{2} & 0 \\ 1 & 0 & 1 \end{pmatrix}. \quad (3)$$

The matrix U_ω is known as ‘magic’ mixing matrix. We will refer to the matrix structures in Eq. (1) as ‘magic’ and ‘cross’ mass matrix, respectively.

There is no observable quark mixing, since $U_{CKM} \equiv U_u^\dagger U_d = U_\omega^\dagger U_\omega = \mathbb{1}_3$. On the contrary, the leptons mix tri-bi-maximally:

$$U_{lepton} \equiv U_e^\dagger U_\nu = U_\omega^\dagger U_\nu = \text{diag}(1, \omega^2, \omega) U_{TBM} \text{diag}(1, 1, -i), \quad U_{TBM} \equiv \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \end{pmatrix}. \quad (4)$$

The diagonal phase matrices can be absorbed by rephasing the charged lepton fields and m_3 . Notice that several different choices of the pair of unitary matrices U_e and U_ν may reproduce TBM mixing. However, as shown in Appendix A, the choice made in Eq. (3) is the only one justified by the A_4 family symmetry.

In ordinary $SO(10)$ models, all light matter fields reside in three dim-16 representations, which also contain right-handed neutrinos. Then, the light Majorana neutrino mass matrix is given in general by

$$m_\nu = M_L - M_\nu M_R^{-1} M_\nu^T, \quad (5)$$

where M_L is a direct Majorana mass term for the left-handed neutrinos, M_ν is the Dirac-type neutrino mass matrix and M_R is the Majorana mass matrix of right-handed neutrinos. The second term in Eq. (5) is the outcome of the type I seesaw mechanism [41, 42, 43, 44, 45], while the first term may arise from a type II seesaw [46, 47, 48, 49]. The latter can be sizable in models with at least one $\overline{126}$ Higgs multiplet and a scalar potential such that the $SU(2)_L$ triplet component of $\overline{126}$ develops a vacuum expectation value (VEV). In all other cases the type I seesaw dominates and then the cross structure of m_ν in Eq. (1) should emerge from the interplay of the M_ν and M_R structures. The simplest possibility is, of course,

$$M_\nu = \begin{pmatrix} A_\nu & 0 & B_\nu \\ 0 & C_\nu & 0 \\ B_\nu & 0 & A_\nu \end{pmatrix}, \quad M_R = \begin{pmatrix} A_R & 0 & B_R \\ 0 & C_R & 0 \\ B_R & 0 & A_R \end{pmatrix}, \quad (6)$$

with at least one among B_ν and B_R being non-zero. Another possibility which may be justified by the A_4 family symmetry is given by

$$M_\nu = \begin{pmatrix} 0 & A_\nu & 0 \\ -A_\nu & 0 & A_\nu \\ 0 & -A_\nu & 0 \end{pmatrix}, \quad (7)$$

with M_R as in Eq. (6). In this case one recovers the form of m_ν in Eq. (1) with $b = -a$, which implies a normal neutrino mass hierarchy.

III. MODELS WITH ONLY RENORMALIZABLE YUKAWA COUPLINGS

In this section we consider the Yukawa couplings $Y\Psi\Psi\Phi$ invariant under $SO(10) \times A_4$ and investigate whether they can generate the mass matrix structures in Eq. (1) and thus realize TBM lepton mixing.

We will indicate with (R, \tilde{R}) a multiplet transforming in the representation R of $SO(10)$ and \tilde{R} of A_4 . Matter fields are unified into a single multiplet $\Psi \sim (16, 3)$. The structure of the mass matrices depends on the assignment of the Higgs multiplets Φ under $SO(10) \times A_4$ and on the components of Φ which acquire a non-zero VEV. We will indicate with $\langle r^\Phi \rangle \equiv v_r^\Phi$ the VEV of the Φ component in the representation r of $SU(5)$. All the possible matrix structures are listed in Appendix B 1.

Let us discuss first the mass matrices of charged fermions. As follows from Eq. (2), the masses of the three families can be accommodated only if the matrix entries A_f , B_f and C_f are all non-zero, of the same order of magnitude, and different from each other. One can generate the up-quark mass matrix M_u introducing the Higgs multiplets $\phi \sim (10, 1)$, $\eta \sim (10, 3)$ and $\rho \sim (120, 3)$. The VEVs

$$\langle 5^\phi \rangle = v_5^\phi, \quad \langle 5_i^\eta \rangle = (v_5^\eta, v_5^\eta, v_5^\eta), \quad \langle 45_i^\rho \rangle = (v_{45}^\rho, v_{45}^\rho, v_{45}^\rho) \quad (8)$$

generate A_u , $B_u + C_u$ and $B_u - C_u$, respectively. Alternatively, ϕ and/or η can be replaced by $\Delta \sim (\overline{126}, 1)$ and/or $\Omega \sim (\overline{126}, 3)$ with the same VEV alignment.

In the down quark and charged lepton sector, there are two classes of contributions to the mass matrices: those of the type $\delta M_e^T = \delta M_d$ can be generated by the VEVs

$$\langle \overline{5}^\phi \rangle = v_5^\phi, \quad \langle \overline{5}_i^\eta \rangle = (v_5^\eta, v_5^\eta, v_5^\eta), \quad \langle \overline{5}_i^\rho \rangle = (v_5^\rho, v_5^\rho, v_5^\rho); \quad (9)$$

those of the type $\delta M_e^T = -3\delta M_d$ can be generated by the VEVs

$$\langle \overline{45}^\Delta \rangle = v_{45}^\Delta, \quad \langle \overline{45}_i^\Omega \rangle = (v_{45}^\Omega, v_{45}^\Omega, v_{45}^\Omega), \quad \langle \overline{45}_i^\rho \rangle = (v_{45}^\rho, v_{45}^\rho, v_{45}^\rho). \quad (10)$$

The first, second and third term in Eqs. (9) and (10) contribute to $A_{d,e}$, $B_{d,e} + C_{d,e}$ and $B_{d,e} - C_{d,e}$, respectively. When all six contributions are present, the three masses of down quarks and charged leptons can be fitted. If some of the VEVs are zero, non-trivial relations between the masses are predicted. For example, an economical scenario with no VEVs in the Higgs doublets of the $\overline{126}$ multiplets would imply $A_e = A_d$ and $B_e + C_e = B_d + C_d$. In this limit one predicts (i) $|e^{i\varphi_\mu} m_\mu + m_\tau| = |e^{i\varphi_s} m_s + m_b|$ with $\varphi_{\mu,s}$ arbitrary, which may be compatible with the GUT scale values (see e.g. [50]), and (ii) $m_e = m_d$, which requires corrections of the order of one MeV.

Let us now move to the neutrino sector. Consider first the term M_L in Eq. (5). It is generated when the $SU(2)_L$ Higgs triplets in $\overline{126}$ multiplets receive a tiny VEV through a type II seesaw mechanism [46, 47, 48, 49]. Taking

$$\langle 15^\Delta \rangle = v_{15}^\Delta, \quad \langle 15_i^\Omega \rangle = (0, v_{15}^\Omega, 0), \quad (11)$$

M_L acquires the structure of m_ν in Eq. (1) with $a = c$. Therefore, if one assumes $m_\nu = M_L$, the TBM lepton mixing is realized with one constraint on the light neutrino mass spectrum [23].

One should stress, however, that the type I seesaw contribution (second term in Eq. (5)) is necessarily present in $SO(10)$ models and it turns out to be incompatible with the TBM structure. In fact, the VEVs v_5^ϕ and v_5^η , which are necessary to generate M_u , also contribute to the neutrino Dirac mass matrix M_ν , which takes the form

$$M_\nu = \begin{pmatrix} A_\nu & B_\nu & B_\nu \\ B_\nu & A_\nu & B_\nu \\ B_\nu & B_\nu & A_\nu \end{pmatrix}, \quad (12)$$

with $A_\nu = A_u$ and $B_\nu = (B_u + C_u)/2$. The right-handed neutrino mass matrix M_R is generated by the VEVs of the $SU(5)$ singlet components of the $\overline{126}$ Higgs multiplets. By constructing the combination $M_\nu M_R M_\nu^T$ for all the possible A_4 structures of M_R (see Table V) and barring fine-tuning of independent couplings, we find that the term $B_\nu \propto v_5^\eta$ is never compatible with the structure of m_ν given in Eq. (1). This shows that the exact realization of the TBM mixing is not possible.

In order to estimate the deviation from TBM mixing, a reasonable hypothesis is that M_R acquires a structure analog to M_L from the VEVs

$$\langle 1^\Delta \rangle = v_1^\Delta, \quad \langle 1_i^\Omega \rangle = (0, v_1^\Omega, 0). \quad (13)$$

In this case M_R has the form in Eq. (6) with $A_R = C_R$ and one finds

$$m_\nu = \begin{pmatrix} a & 0 & b \\ 0 & c & 0 \\ b & 0 & a \end{pmatrix} + \begin{pmatrix} 0 & \epsilon & \epsilon \\ \epsilon & 0 & \epsilon \\ \epsilon & \epsilon & 0 \end{pmatrix}. \quad (14)$$

The effect of type I seesaw is to make $a \neq c$ and to add the ϵ term. It is easy to check that, in the basis in which the charged leptons are diagonal, m_ν is still $\mu - \tau$ symmetric, therefore the values of the mixing angles $\theta_{13} = 0$ and $\theta_{23} = \pi/4$ are preserved. On the contrary, the TBM value of θ_{12} and the light neutrino mass spectrum are corrected in a correlated way. Taking for simplicity all parameters real, one finds

$$\tan 2\theta_{12} = \frac{2\sqrt{2}}{1 + \frac{9\epsilon}{a+b-c}}. \quad (15)$$

The TBM value of θ_{12} is recovered for $\epsilon = 0$. In order to accommodate the experimental value of θ_{12} at the $2\text{-}\sigma$ level [1], one needs $-0.03 < \epsilon/(a+b-c) < 0.04$, so that a few percent type I seesaw contribution can be tolerated. In the case $v_1^\Omega = 0$, the same discussion holds with $a = c$.

We have shown that, if M_u is given by Eq. (1), then the structure of M_ν is not compatible with exact TBM mixing. Vice versa, it is instructive to consider the structures of M_ν leading to TBM mixing and see what are the implications for the up quark sector. A first possibility is to take the M_R structure of Eq. (6), generated by the VEVs in Eq. (13), and the M_ν structure of Eq. (7) by the VEV

$$\langle 5_i^\rho \rangle = (v_5^\rho, 0, v_5^\rho). \quad (16)$$

Even though this VEV alignment does not preserve any subgroup of A_4 , we expect that it may be justified dynamically; the analog alignment has been obtained e.g. in a model with $SU(3)$ family symmetry [51]. This would be sufficient to achieve TBM mixing in the lepton sector. In the context of $SO(10)$, however, the only VEV that may contribute to M_u without modifying the structure of M_ν in Eq. (7) is v_{45}^ρ (see table VI).

	matter	Higgs fields			flavons			
	Ψ	ϕ	ρ	Δ	σ	χ	τ	φ
$SO(10)$	16	10	120	$\overline{126}$	1	1	1	1
A_4	3	1	1	1	1	3	1	3
Z_N	1	α^m	α^m	α^r	α^{-m}	α^{-m}	α^{-r}	α^{-r}

TABLE I: Chiral superfields in a minimal renormalizable model for approximate TBM mixing. Here $\alpha^N = 1$ and $1 \leq m \neq r < N$.

Therefore M_u would be purely antisymmetric ($A_u = 0$ and $B_u = -C_u$) and thus would have one zero and two equal eigenvalues, which clearly is not acceptable. A second possibility to realize exact TBM mixing is to generate the structure of M_ν in Eq. (6) by modifying the VEV alignment of 5_i^η in Eq. (8) as follows:

$$\langle 5_i^\eta \rangle = (0, v_5^\eta, 0). \quad (17)$$

In this case the up quark mass matrix takes the form

$$M_u = \begin{pmatrix} A_u & (B_u - C_u)/2 & C_u \\ (C_u - B_u)/2 & A_u & (B_u - C_u)/2 \\ B_u & (C_u - B_u)/2 & A_u \end{pmatrix}. \quad (18)$$

Since M_d has the structure in Eq. (1), the requirement to have $V_{CKM} \approx \mathbb{1}_3$ would imply $C_u \approx -B_u$ and thus lead to the wrong relation $m_c \approx m_t$ (see Eq.(2)). Vice versa, the requirement to fit the three up quark masses would force the V_{CKM} angles to be large. We conclude that the strong departures from the M_u structure in Eq. (1), required to achieve exact TBM lepton mixing, are not viable phenomenologically.

A comment is in order on the pattern of $SO(10) \times A_4$ spontaneous breaking. The triplet VEVs of the type $(1, 1, 1)$, introduced in Eqs. (8)-(10), break A_4 to a Z_3 subgroup at the electroweak scale. The VEV of the type $(0, 1, 0)$, introduced in Eq. (11), breaks A_4 to a Z_2 subgroup at the scale of light neutrino masses (notice that the large A_4 breaking VEV v_1^Ω in Eq. (13) is not needed in a minimal scenario). This misalignment between the charged fermion sector and the neutrino sector is a crucial ingredient to explain TBM mixing. On the $SO(10)$ side, one faces the problem to generate VEVs only in some specific components of the Higgs multiplets, specifically the Higgs doublets in 10 and 120 multiplets and the Higgs triplets and singlets in $\overline{126}$ multiplets. It is a difficult task to arrange for an appropriate scalar potential, also in view of the different energy scales of different sets of VEVs. The analysis of such potential, which is beyond the scope of this paper, may in principle reveal a connection between the VEV alignment dynamics in the A_4 and $SO(10)$ sectors.

In the rest of this section, we will show that a relatively economic model can be built by using only $SO(10)$ Higgs multiplets which are singlet under A_4 , plus a set of gauge singlet flavon fields responsible for the breaking of the family symmetry. In this approach the problem of VEV alignments can be treated separately in the $SO(10)$ and A_4 sectors. Another advantage is that the number of $SO(10)$ multiplets can be considerably reduced, which is desirable to maintain the theory perturbative well above the GUT scale.

The field content of the model is given in Table I. In order to couple the appropriate flavon fields only to certain $SO(10)$ operators, we introduced a Z_N symmetry, the minimal choice being $N = 3$ with charges $m = 1$ and $r = 2$. The Yukawa superpotential is then given by

$$W_Y = \frac{y_1 \sigma + y_3 \chi}{\Lambda} \Psi \Psi \phi + \frac{g_3 \chi}{\Lambda} \Psi \Psi \rho + \frac{f_1 \tau + f_3 \varphi}{\Lambda} \Psi \Psi \Delta, \quad (19)$$

where Λ is a large energy scale where the $A_4 \times Z_N$ symmetry is realized. The flavon fields are assumed to acquire the VEVs

$$\langle \sigma \rangle = \sigma, \quad \langle \chi_i \rangle = (\chi, \chi, \chi), \quad \langle \tau \rangle = \tau, \quad \langle \varphi_i \rangle = (0, \varphi, 0), \quad (20)$$

which break A_4 as well as Z_N at some scale $\sim \lambda\Lambda$, with $\lambda < 1$. The problem to achieve dynamically such VEV alignment has been addressed and solved in several papers [23, 25, 27, 30]. Notice that in Eq. (19) we included only the leading order operators, linear in the flavon fields. Higher dimensional operators with $1 + n$ flavons generate corrections to the mass matrices of relative order λ^n , which may be significant for small n and λ close to one. However, all the operators up to some given n can be forbidden by using a Z_N with a sufficiently large N and choosing carefully the charges m and r in Table I.

We suppose that all the Higgs doublet in ϕ and ρ acquire a VEV at the electroweak scale, except the one in 5^ρ . Also, 1^Δ acquires a VEV at the GUT scale and the Higgs triplet in 15^Δ takes a VEV of the order of the light neutrino mass scale. Then, the mass matrices of charged fermions are explicitly given by

$$\begin{aligned} M_u &= \frac{1}{\Lambda} \begin{pmatrix} y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi + g_3 \chi v_{45}^\rho & y_3 \chi v_5^\phi - g_3 \chi v_{45}^\rho \\ y_3 \chi v_5^\phi - g_3 \chi v_{45}^\rho & y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi + g_3 \chi v_{45}^\rho \\ y_3 \chi v_5^\phi + g_3 \chi v_{45}^\rho & y_3 \chi v_5^\phi - g_3 \chi v_{45}^\rho & y_1 \sigma v_5^\phi \end{pmatrix}, \\ M_d &= \frac{1}{\Lambda} \begin{pmatrix} y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho + v_{45}^\rho) & y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho + v_{45}^\rho) \\ y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho + v_{45}^\rho) & y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho + v_{45}^\rho) \\ y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho + v_{45}^\rho) & y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho + v_{45}^\rho) & y_1 \sigma v_5^\phi \end{pmatrix}, \\ M_e &= \frac{1}{\Lambda} \begin{pmatrix} y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho - 3v_{45}^\rho) & y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho - 3v_{45}^\rho) \\ y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho - 3v_{45}^\rho) & y_1 \sigma v_5^\phi & y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho - 3v_{45}^\rho) \\ y_3 \chi v_5^\phi - g_3 \chi (v_5^\rho - 3v_{45}^\rho) & y_3 \chi v_5^\phi + g_3 \chi (v_5^\rho - 3v_{45}^\rho) & y_1 \sigma v_5^\phi \end{pmatrix}. \quad (21) \end{aligned}$$

The matrix M_u accommodates the three up quark masses and, combined with M_d , leads to vanishing CKM mixing. The matrices M_d and M_e depend only on four complex parameters, so that non-trivial relations are predicted among their eigenvalues, as discussed after Eq. (10). If one insists to relax this constraint in order to fit independently down quark and charged lepton masses, it suffices to introduce an extra Higgs multiplet $\Delta' \sim (\overline{126}, 1, \alpha^m)$ with a VEV in the $\overline{45}^{\Delta'}$ component. The neutrino mass matrices are given by

$$M_\nu = \frac{1}{\Lambda} \begin{pmatrix} y_1 \sigma & y_3 \chi & y_3 \chi \\ y_3 \chi & y_1 \sigma & y_3 \chi \\ y_3 \chi & y_3 \chi & y_1 \sigma \end{pmatrix} v_5^\phi, \quad M_L = \frac{1}{\Lambda} \begin{pmatrix} f_1 \tau & 0 & f_3 \varphi \\ 0 & f_1 \tau & 0 \\ f_3 \varphi & 0 & f_1 \tau \end{pmatrix} v_{15}^\Delta, \quad M_R = \frac{1}{\Lambda} \begin{pmatrix} f_1 \tau & 0 & f_3 \varphi \\ 0 & f_1 \tau & 0 \\ f_3 \varphi & 0 & f_1 \tau \end{pmatrix} v_1^\Delta. \quad (22)$$

The light neutrino mass matrix has the form in Eq. (14). The departure from the TBM value of the $1-2$ lepton mixing angle is controlled by the ratio $(v_5^\phi)^2 / (v_1^\Delta v_{15}^\Delta)$, where we assume that the couplings in Eq. (19) and also the flavon VEVs in Eq. (20) are all of the same order. As discussed after Eq. (15), this ratio must be smaller than few percents, which is realized e.g. for $v_5^\phi \approx 100$ GeV, $v_1^\Delta \approx 10^{16}$ GeV and $v_{15}^\Delta \approx 0.1$ eV.

In the above model there is no mixing in the quark sector. Let us briefly review possible mechanisms to generate non-vanishing CKM parameters in A_4 models where $V_{CKM} = \mathbb{1}_3$ at leading order. In order to introduce the quark mixing, people considered (i) small explicit A_4 breaking Yukawas [52]; (ii) A_4 breaking in the soft supersymmetry breaking terms [10]; (iii) one-loop corrections that communicate to the quark sector the effect of the A_4 spontaneous breaking in the $(0, 1, 0)$ direction [31, 40]; (iv) spontaneous A_4 breaking by a triplet with VEV (v, v, v_3) with $v \neq v_3$ [35]; (v) extra flavons transforming as $1'$ or $1''$ coupled to the quarks [15, 34]. In most of these cases it is problematic to generate a sufficiently large quark mixing, without introducing too large deviations from TBM lepton mixing [30]. One possible exception is provided by $SU(5)$

GUT models, where the CKM parameters can be introduced as a left-handed rotation in the down quark sector, which corresponds to a right-handed one in the charged lepton sector and therefore does not affect lepton mixing [15]. However, in our analysis of $SO(10)$ GUT models we did not find a contribution to M_d and M_e^T that introduces only a mixing on the left. At the end of section IV we will present a different mechanism to accommodate non-zero CKM parameters, which does not perturb the TBM lepton mixing.

IV. MODELS WITH NON-RENORMALIZABLE OPERATORS

In the previous section we have shown that the TBM lepton mixing cannot be exactly realized by Yukawa couplings symmetric under $SO(10) \times A_4$. In this section we solve this difficulty by considering the effect of non-renormalizable operators that contribute to the fermion mass matrices. In addition, the analysis of such operators will provide a new tool to accommodate the CKM parameters.

Higher dimensional operators are proportional to powers of M_{GUT}/Λ , where the cutoff Λ can be identified with the Planck or the string scale, or with the mass of vector-like matter multiplets. For definiteness, in the following we study this last possibility, by considering the superpotential

$$W = y_A \Psi \Sigma \Phi_A + y_B \Psi \bar{\Sigma} \Phi_B - M \Sigma \bar{\Sigma} . \quad (23)$$

where Φ_A and Φ_B are Higgs multiplets whose components may acquire VEVs a and b , respectively, $(\Sigma, \bar{\Sigma})$ is a vector-like pair of matter multiplets and we assume $a, b \ll M$. In this case Σ and $\bar{\Sigma}$ can be integrated out and one is left with an effective operator

$$W^{eff} = \frac{y_A y_B}{M} (\Psi \Phi_A)_\Sigma (\Psi \Phi_B)_{\bar{\Sigma}} , \quad (24)$$

where the subscripts specify how Ψ and $\Phi_{A,B}$ are contracted. The VEVs a and b can be either of the order of M_{GUT} , if they participate to the $SO(10)$ symmetry breaking to the SM, or of the order of the electroweak scale, if they break the SM gauge group.

The flavour structure generated by the operator in Eq. (24) is determined by the A_4 assignments of Φ_A , Φ_B , Σ and $\bar{\Sigma}$. All the possibilities are analyzed in Appendix B2 and displayed in Table VII. Beside the structures already possible with renormalizable Yukawa couplings, several new flavour structures can be realized (compare with Table V). This provides new options for model-building with A_4 symmetry. Here we will focus on the realization of TBM lepton mixing, but alternative scenarios can be studied on the same footing.

By surveying all the possible operators listed in Table VII, one can identify the flavour structures compatible with the TBM mixing scheme. The cross structure of the neutrino Dirac and Majorana mass matrices, shown in Eqs. (1) and (6), can be built with the following components:

- $diag(1, 1, 1)$ from the operator A;
- $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ from the operator B with VEV alignment $(0, a, 0)$ and no antisymmetric term ($y_A^a = 0$);
- $diag(0, 1, 0)$ from the operator C (or C') with VEV alignment $(0, a, 0)$ and $(0, b, 0)$;
- $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix}$ from the operator C with VEV alignment $(a, 0, a)$ and $(b, 0, b)$;

- $diag(1, 0, 1)$ from the operator D with VEV alignment $(0, a, 0)$ and $(0, b, 0)$ and either $y_{A,B}^a = 0$ or $y_{A,B}^s = 0$;
- $\begin{pmatrix} 1 & 0 & \pm 1 \\ 0 & 2 & 0 \\ \pm 1 & 0 & 1 \end{pmatrix}$ from the operator D with VEV alignment $(a, 0, a)$ and $(b, 0, b)$ and either $y_{A,B}^a = 0$ or $y_{A,B}^s = 0$.

The magic structure of the charged fermion mass matrices, shown in Eq. (1), can be built with the following components:

- $diag(1, 1, 1)$ from the operator A;
- $y_A^s \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} + y_A^a \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}$ from the operator B with VEV alignment (a, a, a) ;
- $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ from the operator C with VEV alignment (a, a, a) and (b, b, b) ;
- $\begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix}$ from the operator C' with VEV alignment (a, a, a) and (b, b, b) ;
- $y_A^s y_B^s \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} + y_A^a y_B^a \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + (y_A^s y_B^a - y_A^a y_B^s) \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}$ from the operator D with VEV alignment (a, a, a) and (b, b, b) .

The contribution of the operator in Eq. (24) to the mass matrices of the different sectors is determined by the $SO(10)$ assignments of Φ_A , Φ_B , Σ and $\bar{\Sigma}$. Higher dimensional operators have been often used to build realistic $SO(10)$ models without large representations, which may be disfavoured theoretically. We thus perform a systematic analysis of dim-5 operators involving only $SO(10)$ multiplets of size less than or equal to 120. There are only six such operators, that are analyzed in Appendix B2 and displayed in Table VIII. Several qualitative new relations between the various sectors are possible, with respect to the case of renormalizable Yukawa couplings (compare with Table VI):

- the operator $16_M 16_M 16_H 16_H$ ($16_M 16_M \overline{16}_H \overline{16}_H$) contributes only to the down (up) sector; here the subscripts distinguish matter (M) from Higgs (H) multiplets;
- the operator $16_M 16_M \overline{16}_H \overline{16}_H$ can generate Majorana masses for left-handed and right-handed neutrinos, with no need of any $\overline{126}_H$ multiplet;
- the operator $(16_M \overline{16}_H)_1 (16_M \overline{16}_H)_1$ contributes to M_ν without affecting M_u ;
- the operator $(16_M 120_H)_{16} (16_M 45_H)_{\overline{16}}$ provides two independent contributions to M_ν and M_u , proportional to $\langle 5_H^{120} \rangle$ and $\langle 45_H^{120} \rangle$, respectively.

We are now in the position to build a minimal model which realizes TBM lepton mixing using the dim-5 operators. Notice first that, in the absence of $\overline{126}_H$ multiplets, the type I seesaw is the dominant source of

light neutrino masses. In fact, the type II contribution to m_ν generated by the operator $16_M 16_M \overline{16}_H \overline{16}_H$ is of the order $\langle 5_{\overline{16}_H} \rangle^2 / M \lesssim (100 \text{ GeV})^2 / M_{GUT} \sim 10^{-3} \text{ eV}$, that is negligible.

In section III we showed that, in the case of type I seesaw, one cannot reproduce TBM lepton mixing by using renormalizable Yukawa couplings. The technical reason is the absence of an operator that provides an off-diagonal symmetric term, needed for the magic structure of M_u , without modifying the cross structure of M_ν . This difficulty can be overcome by using dim-5 operators. The mechanism to generate the required contribution to M_u is most easily described in $SU(5)$ language: the VEV of the up-type doublet in a 45 Higgs multiplet couples (antisymmetrically) to two 10 matter multiplets generating M_u , while it does not contribute to M_ν . In order to make this contribution to M_u not antisymmetric, one needs the insertion of the VEV of a 24 Higgs multiplet, which couples to 10 and $\overline{10}$ matter multiplets. Since the SM singlet in 24 is in the hypercharge direction, the Clebsch-Gordan coefficients for the Q and u^c components are different (by a relative factor -4), thus making M_u not antisymmetric. This mechanism is embedded in $SO(10)$ by the operator $(16_M 120_H)_{16} (16_M 45_H)_{\overline{16}}$, with 45_H^{120} and 24_H^{45} acquiring a VEV. By inspecting Table VIII, this is in fact the only possible contribution to M_u that does not affect M_ν , at least with multiplets of dim ≤ 120 . Notice that such contribution to M_u is symmetric when the VEV of 45_H is in the $B - L$ direction (to see this, replace Eq. (B2) in the row VIII of Table VIII).

For completeness, let us mention another $SU(5)$ mechanism that allows to generate only M_u and not M_ν : one may employ the VEV of a 75 Higgs multiplet, which couples to 10 and $\overline{10}$ (but not to 5 and $\overline{5}$) matter multiplets. The minimal $SO(10)$ embedding is provided by the operator $(16_M 10_H)_{16} (16_M 210_H)_{\overline{16}}$, with 5_H^{10} and 75_H^{210} acquiring a VEV. However, the 75 component that acquires the VEV couples with opposite Clebsch-Gordan coefficients to the Q and u^c components. As a consequence the contribution to M_u of this operator is antisymmetric. A symmetric contribution requires an extra antisymmetric coupling, which can be introduced by replacing 10_H with 120_H .

Our strategy to build an explicit model is to start from the renormalizable model discussed at the end of section III, which is defined by the superpotential in Eq. (19), and replace some of the couplings with the appropriate higher dimensional operators. The field content of the model is given in Table II, where we introduced a Z_N symmetry. The Yukawa superpotential invariant under $SO(10) \times A_4 \times Z_N$ is given by

$$W_Y = \frac{y_1 \sigma}{\Lambda} \Psi \Phi \phi + \frac{g_3 \chi}{\Lambda} \Psi \Psi \rho + \frac{h_3 \chi}{\Lambda} \Psi \overline{\Sigma}_2 A - M_2 \overline{\Sigma}_2 \Sigma_2 + h \Sigma_2 \Psi \rho' + \frac{f_1 \tau + f_3 \varphi}{\Lambda} \Psi \overline{\Sigma}_1 \xi - M_1 \overline{\Sigma}_1 \Sigma_1 + f \Sigma_1 \Psi \xi. \quad (25)$$

In order to forbid all other couplings, up to terms quadratic in the flavons, the Z_N charges in Table II must be carefully chosen. One viable choice is given by $N = 8$, with $m = 1$, $r = 2$ and $n = 5$. By taking N large enough, it is possible to forbid unwanted couplings up to higher order in the flavons. Integrating out the heavy messenger fields $(\Sigma_i, \overline{\Sigma}_i)$, for $i = 1, 2$, the effective superpotential takes the form

$$W_Y^{eff} = \frac{y_1 \sigma}{\Lambda} \Psi \Phi \phi + \frac{g_3 \chi}{\Lambda} \Psi \Psi \rho + \frac{h_3 h \chi}{\Lambda} \frac{(\Psi \rho')_{\Sigma_2} (\Psi A)_{\overline{\Sigma}_2}}{M_2} + \frac{(f_1 \tau + f_3 \varphi) f}{\Lambda} \frac{(\Psi \xi)_{\Sigma_1} (\Psi \xi)_{\overline{\Sigma}_1}}{M_1}. \quad (26)$$

For brevity, in Eqs. (25) and (26) we indicated with h_3 (f_3) two independent couplings h_{3s} and h_{3a} (f_{3s} and f_{3a}), which correspond to the two possible contractions of A_4 indexes in the product $3 \times 3 \times 3$ (see case B in Table VII). The first operator in W_Y^{eff} contributes to the Dirac mass matrices of the four sectors, the second and the third only to the charged fermion mass matrices (we assume $\langle 5^\rho \rangle = \langle 5^{\rho'} \rangle = 0$) and the fourth only to the Dirac and Majorana neutrino mass matrices. The flavons take the same VEVs as in Eq. (20).

The charged fermion mass matrices acquire the magic structure

$$M_f = A_f \mathbb{1}_3 + \frac{B_f + C_f}{2} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} + \frac{B_f - C_f}{2} \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}, \quad (27)$$

	matter fields					Higgs fields					flavons			
	Ψ	Σ_1	$\overline{\Sigma}_1$	Σ_2	$\overline{\Sigma}_2$	ϕ	ρ	ρ'	A	ξ	σ	χ	τ	φ
$SO(10)$	16	1	1	16	$\overline{16}$	10	120	120	45	$\overline{16}$	1	1	1	1
A_4	3	3	3	3	3	1	1	1	1	1	1	3	1	3
Z_N	1	α^{-r}	α^r	α^{-r}	α^r	α^m	α^n	α^r	α^{n-r}	α^r	α^{-m}	α^{-n}	α^{-2r}	α^{-2r}

TABLE II: Chiral superfields of a minimal non-renormalizable model for TBM mixing. Here $\alpha^N = 1$ and $1 \leq m \neq n \neq r < N$.

where the coefficients have the following expression:

$$\begin{aligned}
A_u &= \frac{y_1 \sigma}{\Lambda} v_5^\phi, \\
\frac{B_u + C_u}{2} &= -\frac{h_{3s} h \chi}{\Lambda} \frac{5v_{24}^A}{M_2} v_{45}^{\rho'}, \\
\frac{B_u - C_u}{2} &= \frac{g_3 \chi}{\Lambda} v_{45}^\rho + \frac{h_{3a} h \chi}{\Lambda} \frac{2v_1^A - 3v_{24}^A}{M_2} v_{45}^{\rho'}, \\
A_d &= \frac{y_1 \sigma}{\Lambda} v_5^\phi, \\
\frac{B_d + C_d}{2} &= -\frac{h_{3s} h \chi}{\Lambda} \frac{4v_1^A - v_{24}^A}{M_2} (v_5^{\rho'} + v_{45}^{\rho'}), \\
\frac{B_d - C_d}{2} &= \frac{g_3 \chi}{\Lambda} (v_5^\rho + v_{45}^\rho) - \frac{h_{3a} h \chi}{\Lambda} \frac{2v_1^A - 3v_{24}^A}{M_2} (v_5^{\rho'} + v_{45}^{\rho'}), \\
A_e &= \frac{y_1 \sigma}{\Lambda} v_5^\phi, \\
\frac{B_e + C_e}{2} &= -\frac{h_{3s} h \chi}{\Lambda} \frac{4v_1^A + 9v_{24}^A}{M_2} (v_5^{\rho'} - 3v_{45}^{\rho'}), \\
\frac{B_e - C_e}{2} &= -\frac{g_3 \chi}{\Lambda} (v_5^\rho - 3v_{45}^\rho) + \frac{h_{3a} h \chi}{\Lambda} \frac{2v_1^A - 3v_{24}^A}{M_2} (v_5^{\rho'} - 3v_{45}^{\rho'}). \tag{28}
\end{aligned}$$

These 9 quantities are independent, except for $A_e = A_d$, therefore the 9 masses of quarks and charged leptons can be accommodated with this one constraint, that was already discussed in section III. The hierarchy between the three families requires A_f , $B_f + C_f$ and $B_f - C_f$ to be of the same order. In addition, since the Yukawa of the top is close to one, in the up quark sector the three parameters should be close to the electroweak scale. This implies that the flavon VEVs σ and χ are not much smaller than Λ and the VEV of the adjoint Higgs multiplet, $\langle A \rangle \sim M_{GUT}$, is not much smaller than M_2 . We will discuss in more detail the effect of vector-like matter fields with mass close to M_{GUT} in section V.

The neutrino mass matrices take the form

$$\begin{aligned}
M_\nu &= \frac{1}{\Lambda} \begin{pmatrix} y_1 \sigma v_5^\phi + 2f_{1\tau} \frac{v_1^\xi}{M_1} v_5^\xi & 0 & 2f_{3s} \varphi \frac{v_1^\xi}{M_1} v_5^\xi \\ 0 & y_1 \sigma v_5^\phi + 2f_{1\tau} \frac{v_1^\xi}{M_1} v_5^\xi & 0 \\ 2f_{3s} \varphi \frac{v_1^\xi}{M_1} v_5^\xi & 0 & y_1 \sigma v_5^\phi + 2f_{1\tau} \frac{v_1^\xi}{M_1} v_5^\xi \end{pmatrix}, \\
M_R &= \frac{1}{\Lambda} \begin{pmatrix} f_{1\tau} & 0 & f_{3s} \varphi \\ 0 & f_{1\tau} & 0 \\ f_{3s} \varphi & 0 & f_{1\tau} \end{pmatrix} \frac{(v_1^\xi)^2}{M_1}, \\
M_L &= \frac{1}{\Lambda} \begin{pmatrix} f_{1\tau} & 0 & f_{3s} \varphi \\ 0 & f_{1\tau} & 0 \\ f_{3s} \varphi & 0 & f_{1\tau} \end{pmatrix} \frac{(v_5^\xi)^2}{M_1}. \tag{29}
\end{aligned}$$

Light neutrino masses of the correct order of magnitude require $M_1 \gg v_1^\xi \sim M_{GUT}$. In this case the second term in Eq. (5) can be sufficiently large to accommodate oscillation data. The TBM lepton mixing is exactly realized, as desired.

We conclude this section by showing that the dim-5 operators offer also the opportunity to generate non-zero quark mixing while preserving TBM lepton mixing. The idea is to introduce small deviations from the magic matrix structure in the up quark sector. If these deviations were introduced in the down quark sector, that would affect also the charged lepton mass matrix and, therefore, the TBM values of the lepton mixing angles. We do not explore this possibility in this paper.

Interestingly, the Cabibbo angle is naturally induced if the A_4 triplet VEV aligned in the direction $(0, 1, 0)$, which is needed in the neutrino sector, also contributes to M_u . This can be easily achieved by considering the same model as in Table II, but with Σ_1 and $\bar{\Sigma}_1$ transforming as 45 under $SO(10)$. In this case the fourth operator in Eq. (26) generates the neutrino sector as before (up to irrelevant Clebsch-Gordan factors, see Table VIII), but also contributes to the up quark sector with

$$\delta M_u = \frac{1}{\Lambda} \begin{pmatrix} f_{1\tau} & 0 & f_{3s}\varphi \\ 0 & f_{1\tau} & 0 \\ f_{3s}\varphi & 0 & f_{1\tau} \end{pmatrix} \frac{16v_1^\xi v_5^\xi}{M_1}. \quad (30)$$

We absorb the diagonal entries of δM_u in the definition of A_u in Eq. (28). The off-diagonal entries, instead, do not respect the magic structure. In the basis where M_d is diagonal, the up quark mass matrix is given by

$$U_\omega^T (M_u^{magic} + \delta M_u) U_\omega^* = \begin{pmatrix} m_{u1} + 2\epsilon & -\omega^2\epsilon & -\omega\epsilon \\ -\omega\epsilon & m_{u2} - \epsilon & 2\omega^2\epsilon \\ -\omega^2\epsilon & 2\omega\epsilon & m_{u3} - \epsilon \end{pmatrix}, \quad (31)$$

where m_{ui} are defined in Eq. (2) and $\epsilon \equiv (16/3)(f_{3s}\varphi/\Lambda)(v_1^\xi/M_1)v_5^\xi$. Notice that all entries of M_u receive corrections of the same order. Taking $\epsilon \ll m_{u2,3}$, which is natural since $v_1^\xi/M_1 \ll 1$, one can accommodate the 1 – 2 quark mixing angle, $\theta_{12}^q \approx |\epsilon/m_{u2}| \approx |\epsilon/m_c$. A similar idea to accommodate the Cabibbo angle was used in [40]. The values of the other two quark mixing angles are given by $\theta_{23}^q \sim \theta_{13}^q \sim (m_c/m_t)\theta_{12}^q$, which unfortunately are two small to explain the experimental values.

It is possible to introduce other corrections to M_u in such a way that all CKM parameters can be accommodated, without affecting the structure of the other mass matrices. For this purpose one needs a 120 Higgs multiplet ρ_{up} with VEV only in the $SU(5)$ component 45. Notice that in the model of Table II one can make the identification $\rho \equiv \rho_{up}$, since the VEVs v_5^ρ and v_{45}^ρ are not necessary for the model to work. Then, the operators $\Psi\Psi\rho_{up}$ or $\Psi\Psi A\rho_{up}$ should couple to the flavon φ in order to modify the magic structure of M_u . The flavour structure of such corrections can be sufficiently rich to accommodate all the CKM parameters, e.g. employing the operator D in Table VII. However, we did not find any simple way to explain the hierarchy among the values of the quark mixing angles, therefore we refrain from presenting further details.

V. A MECHANISM TO EXPLAIN THE INTER-FAMILY MASS HIERARCHY

In the previous sections, we analyzed the structure of fermion mixing. We did not address yet the origin of the strong hierarchy between the masses of the three families of quarks and charged leptons. In general, such hierarchy is not explained by the $SO(10) \times A_4$ symmetry by itself. In fact, the mass matrix structure M_f in Eq. (1) accommodates three arbitrary mass eigenvalues, which are therefore free parameters like in the Standard Model. Moreover, as we discussed in the Introduction, since in $SO(10)$ both fermion chiralities

transform as 3 under A_4 , one cannot use an extra family symmetry besides A_4 in order to distinguish the three families.

The purpose of this section is to present an elegant mechanism to generate the hierarchy between the three families in $SO(10) \times A_4$ models. Such mechanism emerges from the analysis of dim-5 operators performed in section IV and it turns out to be closely related to ‘universal seesaw’ models [53, 54, 55, 56]. We will show that it is compatible with the generation of TBM lepton mixing discussed in the previous sections.

The masses of the three families of charged fermions are given in Eq. (2). They are linear combinations, with coefficients of unit modulus, of the mass matrix elements A_f, B_f, C_f . When the equality $A_f = \omega B_f = \omega^2 C_f \equiv m_{f3}/3$ holds, the mass eigenvalues are $(0, 0, m_{f3})$. When $A_f = (m_{f3} + m_{f2})/3$, $B_f = (\omega^2 m_{f3} + \omega m_{f2})/3$ and $C_f = (\omega m_{f3} + \omega^2 m_{f2})/3$, the mass eigenvalues are $(0, m_{f2}, m_{f3})$. If the above relations were approximatively realized, one might explain the inter-family hierarchy. At first sight such relations seem completely *ad hoc* because, in the A_4 models built so far, the parameters A_f, B_f, C_f are generated by independent A_4 invariant operators. To remove this problem, let us begin by writing the mass matrix structure in Eq. (1) as a sum over three rank-1 components:

$$M_f = \frac{m_{f1}}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \frac{m_{f2}}{3} \begin{pmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{pmatrix} + \frac{m_{f3}}{3} \begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix}. \quad (32)$$

The three flavour structures in Eq. (32) cannot be generated in A_4 models with renormalizable Yukawa couplings. However, we have shown in section IV that several new flavour structures are possible when the light matter fields Ψ mix with heavy matter multiplets. This mixing may actually lead to the three requisite structures in Eq. (32). Specifically, the ‘democratic’ structure is obtained from the dim-5 operator in Eq. (24) by taking $(\Sigma, \bar{\Sigma})$ singlets under A_4 and Φ_A and Φ_B transforming as A_4 triplets with VEVs (a, a, a) and (b, b, b) , corresponding to the case C in Table VII. The second (third) flavour structure in Eq. (32) is obtained similarly, but with the pair of matter multiplets $\Sigma'' \sim 1''$ and $\bar{\Sigma}' \sim 1'$ ($\Sigma' \sim 1'$ and $\bar{\Sigma}'' \sim 1''$), corresponding to the case C' in Table VII.

There is a simple group theoretical interpretation of Eq. (32). The A_4 representations decompose under the residual Z_3 symmetry (preserved by the vacuum alignment $(1, 1, 1)$) as $1_{A_4} = 1_{Z_3}$, $1'_{A_4} = 1'_{Z_3}$, $1''_{A_4} = 1''_{Z_3}$ and $3_{A_4} = 1_{Z_3} + 1'_{Z_3} + 1''_{Z_3}$. The mass eigenstates are precisely those three orthogonal combinations of $\Psi_i \sim 3_{A_4}$ which transform in a given representation of Z_3 and, therefore, they separately mix with Σ , Σ' and Σ'' , respectively. The three terms in Eq. (32) correspond to the Z_3 invariants $1_{Z_3} \times 1_{Z_3}$, $1'_{Z_3} \times 1'_{Z_3}$ and $1''_{Z_3} \times 1'_{Z_3}$.

For definiteness, we take the two Higgs VEVs $a \sim M_{EW}$ and $b \sim M_{GUT}$. Since the first and second generation Yukawa couplings are much smaller than one, they are well described by the dim-5 operators obtained decoupling $(\Sigma, \bar{\Sigma})$ and $(\Sigma'', \bar{\Sigma}')$ at the mass scales $M_1 \gg M_2 \gg M_{GUT}$. On the contrary, the third generation Yukawa couplings (in particular the top) are large, therefore we are led to consider a vector-like pair of matter multiplets $(\Sigma', \bar{\Sigma}'')$ with mass M_3 of the same order as M_{GUT} . In this case it is not appropriate to integrate out these states and treat their effect in terms of higher dimensional operators. We shall instead consider explicitly the mixing of Σ' with Ψ and show how this generates the third term in Eq. (32). The ‘inverse’ hierarchy $M_1 \gg M_2 \gg M_3$ may be justified e.g. by a Froggatt-Nielsen $U(1)$ symmetry with different charges for Σ , Σ'' and Σ' .

The above discussion applies to any model with the specified A_4 assignment of fields. In the case of $SO(10)$ models, one should carefully choose the $SO(10)$ operators and symmetry breaking pattern, in order to generate the structure of $M_{u,d,e}$ as in Eq. (32) and preserve, at the same time, the cross structure of m_ν . Let us consider, to begin with, the matter multiplets $\Sigma' \sim (16, 1')$ and $\bar{\Sigma}'' \sim (\bar{16}, 1'')$, together with the

Higgs multiplets $\phi_i \sim (10, 3)$ and $A_i \sim (45, 3)$ (the discussion below can be easily generalized to different $SO(10) \times A_4$ assignments of the fields). The superpotential reads

$$W = -M_3 \Sigma' \bar{\Sigma}'' + y(\Psi_1 \phi_1 + \omega \Psi_2 \phi_2 + \omega^2 \Psi_3 \phi_3) \Sigma' + g(\Psi_1 A_1 + \omega^2 \Psi_2 A_2 + \omega \Psi_3 A_3) \bar{\Sigma}'' . \quad (33)$$

The direct contribution to the light fermion mass matrices from the couplings $\Psi_i \Psi_j \phi_k$ is assumed to be sub-dominant. It can be forbidden e.g. by a parity symmetry Z_2 under which Ψ , ϕ and A are odd.

We assume that A_i acquire VEVs (V, V, V) , such that $\langle A_i \rangle \Psi_{i\alpha} = V k_\alpha \Psi_{i\alpha}$, where α runs over the 16 components of Ψ_i and k_α are Clebsch-Gordan coefficients which depend on the $SO(10)$ direction associated with V . Then, the following linear combination of the 16 multiplet components becomes heavy:

$$\Psi_\alpha^h = c_\alpha \Sigma'_\alpha - \frac{s_\alpha}{\sqrt{3}} (\Psi_{1\alpha} + \omega^2 \Psi_{2\alpha} + \omega \Psi_{3\alpha}) , \quad (34)$$

where

$$c_\alpha \equiv \frac{M_3}{\sqrt{|M_3|^2 + 3|gV k_\alpha|^2}} , \quad s_\alpha \equiv \frac{\sqrt{3}gV k_\alpha}{\sqrt{|M_3|^2 + 3|gV k_\alpha|^2}} . \quad (35)$$

Notice that $|c_\alpha|^2 + |s_\alpha|^2 = 1$. The light fermions $\psi_{i\alpha}^l$, $i = 1, 2, 3$, are given by the three linear combinations orthogonal to Ψ_α^h . The unitary 4×4 mixing matrix is defined by

$$\begin{pmatrix} \Psi_{1\alpha} \\ \Psi_{2\alpha} \\ \Psi_{3\alpha} \\ \Sigma'_\alpha \end{pmatrix} = \begin{pmatrix} & & -s_\alpha^*/\sqrt{3} \\ & P_\alpha & -\omega s_\alpha^*/\sqrt{3} \\ & & -\omega^2 s_\alpha^*/\sqrt{3} \\ Q_\alpha & & c_\alpha^* \end{pmatrix} \begin{pmatrix} \psi_{1\alpha}^l \\ \psi_{2\alpha}^l \\ \psi_{3\alpha}^l \\ \Psi_\alpha^h \end{pmatrix} . \quad (36)$$

Since the choice of basis for $\psi_{i\alpha}^l$ is arbitrary, the matrices P_α and Q_α are determined up to a 3×3 unitary rotation from the right. We choose the basis where $\psi_{i\alpha}^l \rightarrow \Psi_{i\alpha}$ in the limit $s_\alpha \rightarrow 0$. In this case one finds

$$P_\alpha = \mathbb{1}_3 + \frac{|c_\alpha| - 1}{3} \begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix} , \quad Q_\alpha = \frac{s_\alpha c_\alpha^*}{\sqrt{3}|c_\alpha|} \begin{pmatrix} 1 & \omega^2 & \omega \end{pmatrix} . \quad (37)$$

In the following we take c_α real and positive, without loss of generality.

The three light states acquire a mass when the Higgs doublets $H_i^{u,d}$ in ϕ_i acquire VEVs at the electroweak scale. Let us begin from the down quark sector. The second term in Eq. (33) contains the couplings $y(Q_1 H_1^d + \omega Q_2 H_2^d + \omega^2 Q_3 H_3^d) d_{\Sigma'}^c$, as well as $y Q_{\Sigma'} (d_1^c H_1^d + \omega d_2^c H_2^d + \omega^2 d_3^c H_3^d)$. Assuming the VEV alignment (v_d, v_d, v_d) and using Eq. (36) for $\Psi_{i\alpha} = Q_i (d_i^c)$ and $\Sigma'_\alpha = Q_{\Sigma'} (d_{\Sigma'}^c)$, we obtain the following mass term for the three light generations:

$$W \supset (d_1^l \ d_2^l \ d_3^l) \frac{y v_d}{\sqrt{3}} \left[s_Q \begin{pmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{pmatrix} + \begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix} s_{d^c} \right] \begin{pmatrix} d_1^{cl} \\ d_2^{cl} \\ d_3^{cl} \end{pmatrix} . \quad (38)$$

The charged lepton mass term has this same form with the obvious replacements $Q \rightarrow L$ and $d^c \rightarrow e^c$. The two flavour structures in Eq. (38) are exactly those required to reproduce the second and third generation masses as in Eq. (32). The relative hierarchy between the two terms is determined by the direction of the $SO(10)$ -breaking VEV of A . In general $\langle A \rangle = v_3^A T_{3R} + v_{15}^A T_{B-L}$, where T_{3R} (T_{B-L}) is the generator of the right-handed isospin (the $B-L$ symmetry). When $v_{15}^A = 0$, the VEV of A points in the right-handed isospin direction and one has $s_Q = s_L = 0$ as well as $s_{d^c} = s_{e^c}$. In this limit only the third family is massive and $b - \tau$ unification is realized.

Let us now consider the up-quark and Dirac neutrino sectors. If also H_i^u acquire non-zero VEVs (v_u, v_u, v_u) , then M_u (M_ν) takes the same form as in Eq. (38), with the replacements $v_d \rightarrow v_u$ and $d^c \rightarrow u^c$ ($Q \rightarrow L$, $d^c \rightarrow \nu^c$). However, such contribution to M_ν would spoil the TBM lepton mixing. The way to generate only M_u and not M_ν is the same described in section IV: the up-type Higgs doublets which are kept light and acquire a VEV should not be those in $\phi \sim (10, 3)$, but rather those in the $SU(5)$ 45-components of $\rho \sim (120, 3)$. In this case the above derivation can be still applied replacing the coupling $y\Psi\Sigma'\phi$ by $h\Psi\Sigma'\rho$. When $v_{15}^A = 0$, only the top quark acquires a mass.

Next, let us discuss the masses of the second family of charged fermions. One contribution may come from a non-zero $v_{15}^A \ll v_3^A$. In this case $s_Q = -s_L/3 \sim (v_{15}^A/v_3^A)s_{d^c}$, so that the hierarchy between the two terms in Eq. (38) follows from a hierarchy between the $SU(4)_c$ and the $SU(2)_R$ symmetry breaking scales. (A recent Pati-Salam model to obtain a flavour hierarchy from a gauge hierarchy without any family symmetry was recently proposed in Ref. [57].) Notice also that the s and μ masses are generated with the appropriate Georgi-Jarlskog factor, due to their relative $B - L$ number. However, if v_{15}^A contributes also to the up quark sector, one finds $m_c/m_t \sim m_s/m_b$, in strong disagreement with data. The steepest hierarchy in the up quarks cannot be accommodated in this minimal setup.

In fact, there is yet another reason to take the VEV of A in the right-handed isospin direction. We are working in the hypothesis that the cross structure of m_ν in Eq. (1) is generated in the Ψ_i basis. In order to preserve the TBM mixing in the lepton sector, such structure should not be changed by the rotation to the ψ_i^l basis. This is guaranteed by taking $v_{15}^A = 0$, because in this case $s_L = 0$ and therefore the lepton doublets L_i do not mix with $L_{\Sigma'}$.

We set $v_{15}^A = 0$ in the following. In this case, the second family masses may be introduced, as already mentioned, by the mixing with a pair $\Sigma'' \sim (16, 1'')$ and $\bar{\Sigma}' \sim (\bar{16}, 1')$ with mass $M_2 \gg M_3$. One may repeat the analysis done in Eqs. (33)-(38) with some obvious replacements, in particular swapping ω with ω^2 everywhere. Assuming that H_d resides in ϕ and H_u in ρ , consistently with the generation of the third family masses, one can accommodate $m_c/m_t \ll m_s/m_b$ by tuning the independent Yukawa couplings. Some more effort will be needed to make $m_s \neq m_\mu$, as discussed later.

Analogously, one can generate the democratic term in Eq. (32) for the first family masses, by the mixing with a pair $\Sigma \sim (16, 1)$ and $\bar{\Sigma} \sim (\bar{16}, 1)$ with mass $M_1 \gg M_2$. The derivation is again very similar to Eqs. (33)-(38), replacing everywhere ω and ω^2 with 1. Let us remind, however, that since the first family masses are tiny, they may be generated even by non-democratic contributions, from operators close to the Planck scale, which would not affect significantly the values of the mixing angles.

Notice that no CKM mixing between the light quark families is generated by the mechanism described above, as long as the A_4 VEV alignment $(1, 1, 1)$ is preserved in all the operators contributing to M_u and M_d . In this case a Z_3 subgroup is unbroken and, as mentioned above, each of the heavy matter multiplets $\Sigma, \Sigma', \Sigma''$ mixes only with the orthogonal combination of the Ψ_i fields that transforms in the same way under Z_3 . As a consequence, each light family of quarks does not mix with the others. In order to generate the CKM parameters, one should resort to extra contributions to the quark mass matrices which break Z_3 .

Let us implement in an explicit model the mechanism to generate the inter-family mass hierarchy together with TBM lepton mixing and non-zero Cabibbo mixing. In order to correctly describe the masses of third and second family of quarks and charged leptons, we found that the minimal set of multiplets is the one given in Table III. The Yukawa superpotential has the form

$$\begin{aligned}
W_Y = & \Psi \left(\frac{y_3\chi}{\Lambda}\rho_1 + \frac{h_3\chi}{\Lambda}\rho_2 \right) \Sigma' - M_3 \Sigma' \bar{\Sigma}'' + \bar{\Sigma}'' \frac{g_3\chi}{\Lambda} A \Psi \\
& + \Psi \left(\frac{y_2\chi}{\Lambda}\rho_1 + \frac{h_2\chi}{\Lambda}\rho_2 \right) \Sigma'' - M_2 \Sigma'' \bar{\Sigma}' + \bar{\Sigma}' \frac{g_2\chi}{\Lambda} A \Psi .
\end{aligned} \tag{39}$$

	matter fields					Higgs fields			flavon
	Ψ	Σ'	$\bar{\Sigma}''$	Σ''	$\bar{\Sigma}'$	ρ_1	ρ_2	A	χ
	$SO(10)$	16	16	$\bar{16}$	16	$\bar{16}$	120	120	45
A_4	3	1'	1''	1''	1'	1	1	1	3
Z_N	1	α^{-2m}	α^{2m}	α^{-2m}	α^{2m}	α^m	α^m	α^{-3m}	α^m

TABLE III: Chiral superfields needed to realize the inter-family mass hierarchy in the quark and charged lepton sectors. Here $\alpha^N = 1$ and $1 \leq m < N$.

A Z_N symmetry was introduced to forbid all other couplings, up to terms quadratic in the flavons; the minimal choice is $N = 4$ with charge $m = 1$. The adjoint Higgs multiplet A has the VEV v_3^A in the right-handed isospin direction and the triplet flavon has the VEV $\langle \chi_i \rangle = (\chi, \chi, \chi)$. In order to reproduce correctly all the mass ratios, we introduced two 120 multiplets ρ_1 and ρ_2 with the VEVs

$$\langle 45^{\rho_1} \rangle = v_{45}^{\rho_1}, \quad \langle \bar{5}^{\rho_1} \rangle = v_5^{\rho_1}, \quad \langle \bar{45}^{\rho_2} \rangle = v_{45}^{\rho_2}. \quad (40)$$

Following the derivation of Eqs. (33)-(38), it is straightforward to compute the mass eigenvalues defined by Eq. (32):

$$\begin{aligned} m_t &= \frac{\sqrt{3}y_3\chi}{\Lambda} v_{45}^{\rho_1} s_3, & m_c &= \frac{\sqrt{3}y_2\chi}{\Lambda} v_{45}^{\rho_1} s_2, \\ m_b &= - \left(\frac{\sqrt{3}y_3\chi}{\Lambda} v_5^{\rho_1} + \frac{\sqrt{3}h_3\chi}{\Lambda} v_{45}^{\rho_2} \right) s_3, & m_s &= - \left(\frac{\sqrt{3}y_2\chi}{\Lambda} v_5^{\rho_1} + \frac{\sqrt{3}h_2\chi}{\Lambda} v_{45}^{\rho_2} \right) s_2, \\ m_\tau &= \left(\frac{\sqrt{3}y_3\chi}{\Lambda} v_5^{\rho_1} - 3 \frac{\sqrt{3}h_3\chi}{\Lambda} v_{45}^{\rho_2} \right) s_3, & m_\mu &= \left(\frac{\sqrt{3}y_2\chi}{\Lambda} v_5^{\rho_1} - 3 \frac{\sqrt{3}h_2\chi}{\Lambda} v_{45}^{\rho_2} \right) s_2. \end{aligned} \quad (41)$$

where the parameters s_3, s_2 control the mixing between Ψ and Σ', Σ'' and are given by

$$s_3 \equiv \frac{\sqrt{3}g_3\chi v_3^A}{\sqrt{|M_3\Lambda|^2 + |\sqrt{3}g_3\chi v_3^A|^2}}, \quad s_2 \equiv \frac{\sqrt{3}g_2\chi v_3^A}{\sqrt{|M_2\Lambda|^2 + |\sqrt{3}g_2\chi v_3^A|^2}}. \quad (42)$$

The heaviness of the top requires to take $M_3 \sim v_3^A$ so that $s_3 \sim 1$. The hierarchy between second and third generation masses is then explained by taking $M_2 \gg v_3^A$, so that $s_2 \ll 1$. Approximate $b - \tau$ unification is realized when the first term in m_b and m_τ dominates over the second. The ratio $m_s/m_\mu \sim 1/3$ is realized when the second term in m_s and m_μ dominates over the first. More precisely, the six masses in Eq. (41) are constrained by one non-trivial relation, $(3m_s - m_\mu) = (m_c/m_t)(3m_b - m_\tau)$, which connects the two phenomenological facts $m_s/m_b \gg m_c/m_t$ and $m_\mu \approx 3m_s$ at the GUT scale. At this level the first family masses are vanishing.

As for the neutrino sector, we need to generate the cross structure of m_ν , as required by TBM mixing. One may think that is sufficient to introduce the last operator in Eq. (26), however the size of neutrino masses is too small in this case, as can be seen by inspecting Eq. (29) (in the present scenario the term $y_1\sigma v_5^\phi$ in M_ν is absent). To solve this problem, the neutrino Dirac mass matrix M_ν and the right-handed neutrino Majorana mass matrix M_R should be generated by two independent operators. We found that the minimal set of multiplets to achieve this purpose is the one given in Table IV, with two $\bar{16}$ Higgs multiplets taking VEVs only in the directions $\langle 1^{\xi_M} \rangle = v_1^{\xi_M}$ and $\langle 5^{\xi_D} \rangle = v_5^{\xi_D}$. We introduced an auxiliary symmetry $Z_{N'}$ such that all fields charged under Z_N are neutral under $Z_{N'}$ and vice versa. Then, the superpotential in Eq. (39) is extended to include the extra terms

$$W_Y \supset \Psi \left(\frac{f_{1M\tau} + f_{3M\varphi}}{\Lambda} \xi_M + \frac{f'_{\sigma\sigma}}{\Lambda} \xi_D \right) \bar{\Sigma}_M - M_M \bar{\Sigma}_M \Sigma_M + f_M \Sigma_M \xi_M \Psi$$

	matter fields					Higgs fields		flavons		
	Ψ	Σ_D	$\bar{\Sigma}_D$	$\bar{\Sigma}_M$	Σ_M	ξ_D	ξ_M	σ	τ	φ
$SO(10)$	16	1	1	45	45	$\bar{16}$	$\bar{16}$	1	1	1
A_4	3	3	3	3	3	1	1	1	1	3
$Z_{N'}$	1	α^{-n}	α^n	α^{-r}	α^r	α^n	α^{-r}	α^{r-n}	α^{2r}	α^{2r}

TABLE IV: Chiral superfields needed to generate the cross structure of the neutrino mass matrix and the Cabibbo mixing angle. Here $\alpha^{N'} = 1$ and $1 \leq n \neq r < N'$.

$$+ \Psi \frac{f_{\sigma\sigma}}{\Lambda} \xi_M \bar{\Sigma}_D - M_D \Sigma_D \bar{\Sigma}_D + f_D \Sigma_D \xi_D \Psi . \quad (43)$$

The $Z_{N'}$ assignments of Table IV forbid all other $SO(10) \times A_4$ invariant couplings (a minimal viable choice is $N' = 5$, with charges $n = 2$ and $r = 4$). Once the messenger fields $(\Sigma_D, \bar{\Sigma}_D)$ and $(\Sigma_M, \bar{\Sigma}_M)$ are integrated out, the effective superpotential contains two relevant operators:

$$W_Y^{eff} \supset \frac{(f_{1M}\tau + f_{3M}\varphi)f_M}{\Lambda} \frac{(\Psi\xi_M)_{\bar{\Sigma}_M}(\Psi\xi_M)_{\Sigma_M}}{M_M} + \frac{f_{\sigma}f_D\sigma}{\Lambda} \frac{(\Psi\xi_M)_{\bar{\Sigma}_D}(\Psi\xi_D)_{\Sigma_D}}{M_D} . \quad (44)$$

The first operator generates the cross structure of M_R , while the second generates $M_\nu \propto \mathbb{1}_3$ (remember that only the singlet in ξ_M and the doublet in ξ_D acquire a non-zero VEV). By taking $M_M \gg M_D \geq M_{GUT}$, one obtains right-handed neutrino masses significantly below M_{GUT} and keeps the Dirac neutrino masses close to electroweak scale. In this way sufficiently large neutrino masses may be generated by type I seesaw.

When also $\langle 5^{\xi_M} \rangle = v_5^{\xi_M}$ is different from zero, the first operator in Eq. (44) gives a negligible contribution to M_ν , because $M_M \gg M_D$. However, it plays an important role in the up quark sector, since it provides a small correction to M_u in a form completely analog to Eq. (30). This correction generates the Cabibbo mixing angle, in the same way as discussed at the end of section IV.

In summary, the model defined by the superpotential in Eqs. (39) and (43) accounts for TBM mixing in the lepton sector and the 1 – 2 mixing in the quark sector, as well as for the hierarchical values of the quark and charged lepton masses.

Before concluding, let us remark that, in most models of ‘universal seesaw’ with a realistic phenomenology, the mass terms $M\Sigma\bar{\Sigma}$ of the messenger fields are sensitive to the $SO(10)$ breaking VEVs, thus giving different masses to the different components of Σ . One may also introduce a non-trivial mixing between the heavy matter families. In our construction we barred these extra possibilities for simplicity. A recent model of ‘universal seesaw’ making use of rank-1 flavour structures, together with more references, can be found in [58]. Heavy messenger fields are also employed in a similar fashion in an $SO(10) \times SU(3)$ model for TBM mixing [21].

VI. CONCLUSIONS

In this paper we performed a systematic analysis of fermion mass matrices in models of Grand Unification based on the gauge symmetry $SO(10)$ with the discrete family symmetry A_4 . In these models all light fermions and right-handed neutrinos may be unified in a single multiplet $(16, 3)$ of the group $SO(10) \times A_4$. We demonstrated that, even though this scenario is very constrained, it is possible to understand the disparity between the quark and lepton mixing angles as well as the strong hierarchy between the three families of charge fermion masses.

In models with renormalizable Yukawa couplings, we found that the exact TBM lepton mixing can be obtained only if the type I seesaw contribution to m_ν is neglected. The specific effect of such contribution is

to shift the $1 - 2$ lepton mixing angle from the tri-maximal value. The non-zero values of the CKM mixing angles cannot be accommodated without introducing further departures from TBM lepton mixing.

A much richer flavour structure appears once the effect of higher dimensional operators is taken into account. We considered dim-5 operators generated by the mixing of light families with heavy vector-like matter multiplets: the corresponding structures of the mass matrices are listed in Appendix B 2, for all A_4 representations and all the $SO(10)$ representations of size ≤ 120 . This classification proves to be a useful tool for model-building.

We found that the dim-5 operators help to evade several difficulties in the construction of a satisfactory $SO(10) \times A_4$ model of flavour. One crucial ingredient is provided by those operators which contribute differently to the Dirac mass matrices M_u and M_ν . First, it is possible to obtain exact TBM lepton mixing from a type I seesaw. Second, there are contributions which generate non-zero CKM mixing angles without disturbing the TBM pattern in the lepton sector.

Moreover, we have shown that the mixing of the three light families with vector-like matter multiplets provides a natural explanation of the inter-family mass hierarchy of quarks and charged leptons. In fact, rank-1 contributions to the mass matrices are generated by the mixing with heavy dim-16 multiplets, that transform in a dim-1 representation of A_4 . The flavour structures of such contributions are exactly those necessary to achieve TBM mixing. A hierarchy in the masses of the heavy families and/or in the VEVs breaking $SO(10)$ reflects into a hierarchy of the masses of the light families.

Acknowledgments

We thank José F. W. Valle for useful comments and encouragement. MF thanks the Aspen Center for Physics (2007 Workshop on “Neutrino physics: looking forward”), where this project was conceived by discussing with Ernest Ma and José F. W. Valle, and the Instituto de Física Corpuscular of Valencia, for the kind hospitality. The work of FB and SM was supported by MEC grant FPA2005-01269 and FPA2008-01935-E, by EC RTN network MRTN-CT-2004-503369, and by Generalitat Valenciana ACOMP06/154. MF was supported in part by the Marie Curie Intra-European Fellowship MEIF-CT-2007-039968, the CNRS/USA exchange grant 3503, and the RTN European Program MRTN-CT-2004-503369.

APPENDIX A: THE GROUP A_4 AND ITS BREAKING PATTERN

The pair of lepton mass matrix structures defined in in Eq. (1) leads to exact TBM mixing in the lepton sector. In this Appendix we prove that such realization of TBM mixing is the only one that can be obtained by the spontaneous breaking of an A_4 flavour symmetry. We follow an approach already applied to the group A_4 in Ref. [31] and to the group T' in Ref. [59] (see also [60]).

The discrete group A_4 is formed by the even permutations of four objects. It can be defined by two generators S and T such that

$$S^2 = T^3 = (ST)^3 = 1. \quad (\text{A1})$$

The 12 elements of A_4 belong to four conjugacy classes

$$\begin{aligned} \mathcal{C}_1 &: I ; \\ \mathcal{C}_2 &: T, ST, TS, STS ; \\ \mathcal{C}_3 &: T^2, ST^2, T^2S, TST ; \\ \mathcal{C}_4 &: S, T^2ST, TST^2 . \end{aligned} \quad (\text{A2})$$

Each element g of $\mathcal{C}_{2,3}$ (\mathcal{C}_4) satisfies $g^3 = 1$ ($g^2 = 1$) and, therefore, generates a subgroup Z_3 (Z_2) of A_4 . There are three dim-1 and one dim-3 irreducible representations. We adopt the basis where the generators of the dim-3 representation are given by

$$S = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}. \quad (\text{A3})$$

We assume that both lepton doublets and singlets transform as A_4 triplets, as dictated by $SO(10)$. The lepton mixing matrix $U_{lepton} = U_e^\dagger U_\nu$ depends on how A_4 is spontaneously broken in the charged lepton and in the neutrino sector. If A_4 were unbroken in one sector, the mass matrix would be proportional to the identity. If A_4 were broken completely in one sector, the mass matrix would be generic and the values of the mixing angles arbitrary. Therefore, the only non-trivial case occurs when A_4 is broken to a subgroup in the charged lepton sector and to another subgroup in the neutrino sector. There are three possibilities:

- the VEV of an A_4 triplet with alignment $(1, 1, 1)$ breaks A_4 to Z_3 ;
- the VEV of an A_4 triplet with alignment $(0, 1, 0)$ breaks A_4 to Z_2 ;
- the VEV of a singlet in the $1'$ or $1''$ representation breaks A_4 to $Z_2 \times Z_2$.

The relevant choice to obtain TBM mixing is to preserve Z_3 in one sector and Z_2 in the other, as we now show (one may check in a similar way that all other choices do not lead to TBM mixing).

If g is the generator of a subgroup of A_4 , the mass term $\psi M \psi^c$ is invariant with respect to g if and only if $g M g^T = M$. For definiteness, and without loss of generality, let us consider the Z_3 and Z_2 subgroups generated by T and S , respectively. There are two possibilities

$$\begin{aligned} (i) \quad T M_e T^T = M_e &\Rightarrow M_e = \begin{pmatrix} A_e & B_e & C_e \\ C_e & A_e & B_e \\ B_e & C_e & A_e \end{pmatrix}; \quad S m_\nu S^T = m_\nu &\Rightarrow m_\nu = \begin{pmatrix} a & 0 & b \\ 0 & c & 0 \\ b & 0 & d \end{pmatrix}; \\ (ii) \quad S M_e S^T = M_e &\Rightarrow M_e = \begin{pmatrix} A_e & 0 & C_e \\ 0 & D_e & 0 \\ B_e & 0 & E_e \end{pmatrix}; \quad T m_\nu T^T = m_\nu &\Rightarrow m_\nu = \begin{pmatrix} a & b & b \\ b & a & b \\ b & b & a \end{pmatrix}. \end{aligned} \quad (\text{A4})$$

In the case (i) we recognize the charged lepton mass matrix M_e of Eq. (1). Notice that m_ν is more general than in Eq. (1), because the Z_2 invariance allows different entries on the diagonal, in particular $a \neq d$. This is because one may couple to the neutrinos also a Higgs transforming as $1'$ or $1''$, and still maintain m_ν invariant under Z_2 [31]. Therefore this A_4 breaking pattern is not sufficient to predict TBM mixing. If one does not introduce such $1'$ and $1''$ fields in the model, one finds $a = c = d$ and the neutrino mass matrix of Eq. (1) is recovered in the special limit $a = c$. Then exact TBM lepton mixing is obtained.

For what concerns the case (ii), one can obtain TBM mixing in the neutrino sector alone, by taking M_e diagonal. This scenario is realized by coupling to the charged leptons three Higgs singlets $1, 1', 1''$ and no triplet (the A_4 subgroup $Z_2 \times Z_2$ is preserved in this case). However, even though m_ν is diagonalized by U_{TBM} , one has $m_1 = m_3$, in strong disagreement with oscillation experiments. Therefore the case (ii) is not viable.

case	A_4 operator	mass matrix structure
A	$3_M 3_M 1_H$	$y \text{diag}(1, 1, 1) v$
A'	$3_M 3_M 1'_H$	$y' \text{diag}(1, \omega, \omega^2) v'$
A''	$3_M 3_M 1''_H$	$y'' \text{diag}(1, \omega^2, \omega) v''$
B	$3_M 3_M 3_H$	$y^s \begin{pmatrix} 0 & v_3 & v_2 \\ v_3 & 0 & v_1 \\ v_2 & v_1 & 0 \end{pmatrix} + y^a \begin{pmatrix} 0 & v_3 & -v_2 \\ -v_3 & 0 & v_1 \\ v_2 & -v_1 & 0 \end{pmatrix}$

TABLE V: The possible A_4 flavour structures arising from renormalizable Yukawa couplings. The subscript M (H) identifies the matter (Higgs) multiplets.

case	$SO(10)$ operator	mass matrices
I	$16_M 16_M 10_H$	$M_u = M_\nu = Y_{10} v_5$ $M_d = M_e = Y_{10} v_{\overline{5}}$
II	$16_M 16_M 120_H$	$M_u = Y_{120} v_{45}$ $M_\nu = Y_{120} v_5$ $M_d = Y_{120} (v_{\overline{5}} + v_{\overline{45}})$ $M_e^T = Y_{120} (v_{\overline{5}} - 3v_{\overline{45}})$
III	$16_M 16_M \overline{126}_H$	$M_u = Y_{\overline{126}} v_5$ $M_\nu = -3Y_{\overline{126}} v_5$ $M_d = Y_{\overline{126}} v_{\overline{45}}$ $M_e = -3Y_{\overline{126}} v_{\overline{45}}$ $M_L = Y_{\overline{126}} v_{15}$ $M_R = Y_{\overline{126}} v_1$

TABLE VI: The contributions to the mass matrices from $SO(10)$ invariant renormalizable Yukawa couplings. Different VEVs of the same $SO(10)$ Higgs multiplet carry a subscript indicating the $SU(5)$ component they belong to.

APPENDIX B: STRUCTURES OF MASS MATRICES WITH $SO(10) \times A_4$ SYMMETRY

1. Renormalizable Yukawa couplings

Consider the Yukawa couplings $Y_{ij} \Psi_i \Psi_j \Phi$, where the matter multiplets transform as $\Psi \sim (16, 3)$ under $SO(10) \times A_4$ and Φ is a Higgs multiplet. For all possible $SO(10) \times A_4$ assignments of Φ , let us analyze the structure of the quark and lepton mass matrices that are generated when a given component of Φ acquires a VEV v .

The A_4 assignment of Φ determines the flavour structure of the mass matrices, as shown in Table V. When $\Phi \sim 3$ (case B in the Table), there are two A_4 -invariants, one symmetric and the other antisymmetric in the flavour indexes. The $SO(10)$ assignment of Φ determines the relative contribution to the mass matrices of the different sectors, as shown in Table VI. The Yukawa coupling matrices Y_{10} and $Y_{\overline{126}}$ (Y_{120}) are (anti)symmetric in the flavour indexes.

For a given $SO(10) \times A_4$ assignment of Φ , the contribution to the mass matrices is obtained combining the corresponding rows in Tables V and VI. Such contribution is non-zero only if both rows contain terms (anti)symmetric in the flavour indexes.

2. Non-renormalizable operators

Consider the generic dimension-5 operator $c_{ij}\Psi_i\Psi_j\Phi_A\Phi_B/M$, where $\Phi_{A,B}$ are two (possibly equal) Higgs multiplets and M is a cutoff scale. When the appropriate components of Φ_A and Φ_B acquire VEVs a and b , respectively, one generates contributions to the quark and lepton mass matrices. Such operator may arise from the exchange of a pair of vector-like matter multiplets $(\Sigma, \bar{\Sigma})$ of mass M , with the superpotential given in Eq. (23). If one is interested in physics at scales much smaller than M (that is, for $a, b \ll M$), the pair $(\Sigma, \bar{\Sigma})$ can be integrated out and one is left with the effective superpotential

$$W^{eff} = \frac{y_A y_B}{M} (\Psi\Phi_A)_\Sigma (\Psi\Phi_B)_{\bar{\Sigma}}. \quad (\text{B1})$$

The dim-5 operators may also arise from the exchange of heavy Higgs multiplets Φ' . We do not consider this possibility, since in this case the operators involve the Yukawa coupling $Y_{ij}\Psi_i\Psi_j\Phi'$, therefore they do not generate new mass matrix structures, besides those already discussed in Appendix B 1.

The A_4 assignments of $\Phi_{A,B}$ and $(\Sigma, \bar{\Sigma})$ determine the flavour structure of the mass matrices. There are 9 possible cases listed in Table VII. The 5 cases B', B'', C, C', D provide new flavour structures with respect to the Yukawa couplings in Table V. Let us illustrate some new possibilities:

- In the case C (and C') each entry of the mass matrix is proportional to a different pair of VEVs, therefore matrices with only one non-zero entry (row, column) can be generated choosing the appropriate VEV alignment.
- In the case D, by taking the alignment $(0, a, 0)$ and $(0, b, 0)$ only the entries (11) and (33) are generated, with relative coefficients $(y_A^s \pm y_A^a)(y_B^s \pm y_B^a)$.
- In the case D, by taking the alignment (a, a, a) and $(b, 0, 0)$, only the entries 12, 13, 22, 33 are generated; if $y_{A,B}^a = 0$, the 4 entries are equal.

The $SO(10)$ assignments of $\Phi_{A,B}$ and $(\Sigma, \bar{\Sigma})$ determine the relative contribution to the mass matrices of the different sectors. In Table VIII we classify all dim-5 operators involving representations with size ≤ 120 . It is useful to recall that an $SO(10)$ adjoint Higgs multiplet 45_H can acquire VEVs in two independent directions. In the Table we indicate with b_1 (b_{24}) the VEV in the $SU(5)$ singlet (adjoint) direction. Equivalently, one can write the VEV of 45_H as a combination of a VEV b_3 in the right-handed isospin direction and a VEV b_{15} in the $B - L$ direction, by using

$$b_1 = \frac{1}{5}(b_3 + 3b_{15}), \quad b_{24} = \frac{1}{5}(-b_3 + 2b_{15}). \quad (\text{B2})$$

The Clebsch-Gordan coefficients in Table VIII can be derived e.g. with the help of Refs.[61, 62, 63, 64, 65].

For a given $SO(10) \times A_4$ assignment of Φ_A, Φ_B, Σ and $\bar{\Sigma}$, the contribution to the mass matrices is obtained by combining the corresponding rows of Tables VII and VIII.

Operators with dimension larger than 5 can also correct significantly fermion masses, in particular when the cutoff M is not much larger than M_{GUT} . An $SO(10) \times A_4$ model which crucially relies on a dim-6 operator was built in [19]. In this model the operator VIII of Table VIII is used, with 120_H replaced by a product $45_H 10_H$.

case	A_4 operator	mass matrix structure
A	$(3_M 1_H)_3 (3_M 1_H)_3$ $(3_M 1'_H)_3 (3_M 1''_H)_3$	$y_{AYB} \text{diag}(1, 1, 1) \frac{ab}{M}$
A'	$(3_M 1_H)_3 (3_M 1'_H)_3$ $(3_M 1''_H)_3 (3_M 1''_H)_3$	$y'_{AY'B} \text{diag}(1, \omega, \omega^2) \frac{ab}{M}$
A''	$(3_M 1_H)_3 (3_M 1''_H)_3$ $(3_M 1'_H)_3 (3_M 1'_H)_3$	$y''_{AY''B} \text{diag}(1, \omega^2, \omega) \frac{ab}{M}$
B	$(3_M 3_H)_3 (3_M 1_H)_3$	$\left[y_{AYB}^s \begin{pmatrix} 0 & a_3 & a_2 \\ a_3 & 0 & a_1 \\ a_2 & a_1 & 0 \end{pmatrix} + y_{AYB}^a \begin{pmatrix} 0 & a_3 & -a_2 \\ -a_3 & 0 & a_1 \\ a_2 & -a_1 & 0 \end{pmatrix} \right] \frac{b}{M}$
B'	$(3_M 3_H)_3 (3_M 1'_H)_3$	$\left[y_{AYB}^s \begin{pmatrix} 0 & a_3 \omega & a_2 \omega^2 \\ a_3 & 0 & a_1 \omega^2 \\ a_2 & a_1 \omega & 0 \end{pmatrix} + y_{AYB}^a \begin{pmatrix} 0 & a_3 \omega & -a_2 \omega^2 \\ -a_3 & 0 & a_1 \omega^2 \\ a_2 & -a_1 \omega & 0 \end{pmatrix} \right] \frac{b}{M}$
B''	$(3_M 3_H)_3 (3_M 1''_H)_3$	$\left[y_{AYB}^s \begin{pmatrix} 0 & a_3 \omega^2 & a_2 \omega \\ a_3 & 0 & a_1 \omega \\ a_2 & a_1 \omega^2 & 0 \end{pmatrix} + y_{AYB}^a \begin{pmatrix} 0 & a_3 \omega^2 & -a_2 \omega \\ -a_3 & 0 & a_1 \omega \\ a_2 & -a_1 \omega^2 & 0 \end{pmatrix} \right] \frac{b}{M}$
C	$(3_M 3_H)_1 (3_M 3_H)_1$	$y_{AYB} \begin{pmatrix} a_1 b_1 & a_1 b_2 & a_1 b_3 \\ a_2 b_1 & a_2 b_2 & a_2 b_3 \\ a_3 b_1 & a_3 b_2 & a_3 b_3 \end{pmatrix} \frac{1}{M}$
C'	$(3_M 3_H)_{1'} (3_M 3_H)_{1''}$	$y_{AYB} \begin{pmatrix} a_1 b_1 & a_1 b_2 \omega^2 & a_1 b_3 \omega \\ a_2 b_1 \omega & a_2 b_2 & a_2 b_3 \omega^2 \\ a_3 b_1 \omega^2 & a_3 b_2 \omega & a_3 b_3 \end{pmatrix} \frac{1}{M}$
D	$(3_M 3_H)_3 (3_M 3_H)_3$	$\left[y_{AYB}^s \begin{pmatrix} a_2 b_2 + a_3 b_3 & a_2 b_1 & a_3 b_1 \\ a_1 b_2 & a_3 b_3 + a_1 b_1 & a_3 b_2 \\ a_1 b_3 & a_2 b_3 & a_1 b_1 + a_2 b_2 \end{pmatrix} \right. \\ \left. + y_{AYB}^a \begin{pmatrix} a_2 b_2 + a_3 b_3 & -a_2 b_1 & -a_3 b_1 \\ -a_1 b_2 & a_3 b_3 + a_1 b_1 & -a_3 b_2 \\ -a_1 b_3 & -a_2 b_3 & a_1 b_1 + a_2 b_2 \end{pmatrix} \right. \\ \left. + y_{AYB}^s \begin{pmatrix} -a_2 b_2 + a_3 b_3 & -a_2 b_1 & a_3 b_1 \\ a_1 b_2 & -a_3 b_3 + a_1 b_1 & -a_3 b_2 \\ -a_1 b_3 & a_2 b_3 & -a_1 b_1 + a_2 b_2 \end{pmatrix} \right. \\ \left. + y_{AYB}^a \begin{pmatrix} -a_2 b_2 + a_3 b_3 & a_2 b_1 & -a_3 b_1 \\ -a_1 b_2 & -a_3 b_3 + a_1 b_1 & a_3 b_2 \\ a_1 b_3 & -a_2 b_3 & -a_1 b_1 + a_2 b_2 \end{pmatrix} \right] \frac{1}{M}$

TABLE VII: The A_4 flavour structures arising from dim-5 operators, defined as in Eq. (B1). In the cases A, A', A'', C and D one may identify both the two Higgs multiplets, $\Phi_A \equiv \Phi_B$, and the pair of messengers, $\Sigma \equiv \bar{\Sigma}$. If these identifications are made, one has to take $y_A \equiv y_B$ (with possible superscripts understood).

-
- [1] M. Maltoni, T. Schwetz, M. A. Tortola, and J. W. F. Valle, New J. Phys. **6**, 122 (2004), hep-ph/0405172 v6.
[2] G. L. Fogli, E. Lisi, A. Marrone, and A. Palazzo, Prog. Part. Nucl. Phys. **57**, 742 (2006), hep-ph/0506083.
[3] C. H. Albright, K. S. Babu, and S. M. Barr, Phys. Rev. Lett. **81**, 1167 (1998), hep-ph/9802314.
[4] K. S. Babu, J. C. Pati, and F. Wilczek, Nucl. Phys. **B566**, 33 (2000), hep-ph/9812538.
[5] B. Bajc, G. Senjanovic, and F. Vissani, Phys. Rev. Lett. **90**, 051802 (2003), hep-ph/0210207.
[6] H. S. Goh, R. N. Mohapatra, and S.-P. Ng, Phys. Lett. **B570**, 215 (2003), hep-ph/0303055.

case	$SO(10)$ operator	mass matrices
IV	$(16_M 16_H)_{10} (16_M 16_H)_{10}$	$M_d = K a_{\bar{5}} b_1 + K^T a_1 b_{\bar{5}}$ $M_e^T = K a_{\bar{5}} b_1 + K^T a_1 b_{\bar{5}}$
V	$(16_M \overline{16}_H)_1 (16_M \overline{16}_H)_1$	$M_\nu = K a_5 b_1 + K^T a_1 b_5$ $M_L = K_s a_5 b_5$ $M_R = K_s a_1 b_1$
VI	$(16_M \overline{16}_H)_{45} (16_M \overline{16}_H)_{45}$	$M_u = 8K_s (a_5 b_1 + a_1 b_5)$ $M_\nu = 3(K a_5 b_1 + K^T a_1 b_5)$ $M_L = -5K_s a_5 b_5$ $M_R = -5K_s a_1 b_1$
VII	$(16_M 10_H)_{16} (16_M 45_H)_{\overline{16}}$	$M_u = K a_5 (b_1 - 4b_{24}) + K^T a_5 (b_1 + b_{24})$ $M_\nu = 5K a_5 b_1 + K^T a_5 (-3b_1 - 3b_{24})$ $M_d = K a_{\bar{5}} (-3b_1 + 2b_{24}) + K^T a_{\bar{5}} (b_1 + b_{24})$ $M_e^T = K a_{\bar{5}} (-3b_1 - 3b_{24}) + K^T a_{\bar{5}} (b_1 + 6b_{24})$
VIII	$(16_M 120_H)_{16} (16_M 45_H)_{\overline{16}}$	$M_u = K a_{45} (b_1 - 4b_{24}) - K^T a_{45} (b_1 + b_{24})$ $M_\nu = 5K a_5 b_1 - K^T a_5 (-3b_1 - 3b_{24})$ $M_d = K (a_{\bar{5}} + a_{\overline{45}}) (-3b_1 + 2b_{24}) - K^T (a_{\bar{5}} + a_{\overline{45}}) (b_1 + b_{24})$ $M_e^T = K (a_{\bar{5}} - 3a_{\overline{45}}) (-3b_1 - 3b_{24}) - K^T (a_{\bar{5}} - 3a_{\overline{45}}) (b_1 + 6b_{24})$
IX	$(16_M 16_H)_{120} (16_M 16_H)_{120}$	$M_d = K (a_{\bar{5}} b_1 + 2a_1 b_{\bar{5}}) + K^T (a_1 b_{\bar{5}} + 2a_{\bar{5}} b_1)$ $M_e^T = K (a_{\bar{5}} b_1 + 2a_1 b_{\bar{5}}) + K^T (a_1 b_{\bar{5}} + 2a_{\bar{5}} b_1)$

TABLE VIII: The contributions to the mass matrices from $SO(10)$ -invariant dim-5 operators, defined as in Eq. (B1). Here we introduced $K \equiv y_A y_B / M$ and $K_s \equiv (K + K^T) / 2$. This list exhausts all possibilities with Φ_A, Φ_B, Σ and $\overline{\Sigma}$ in representations with size ≤ 120 . Different VEVs of the same $SO(10)$ Higgs multiplet carry a subscript indicating the $SU(5)$ component they belong to.

- [7] K. S. Babu and S. M. Barr, Phys. Lett. **B381**, 202 (1996), hep-ph/9511446.
- [8] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B530**, 167 (2002), hep-ph/0202074.
- [9] E. Ma and G. Rajasekaran, Phys. Rev. **D64**, 113012 (2001), hep-ph/0106291.
- [10] K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. **B552**, 207 (2003), hep-ph/0206292.
- [11] D. Wyler, Phys. Rev. **D19**, 3369 (1979).
- [12] G. C. Branco, H. P. Nilles, and V. Rittenberg, Phys. Rev. **D21**, 3417 (1980).
- [13] E. Ma, H. Sawanaka, and M. Tanimoto, Phys. Lett. **B641**, 301 (2006), hep-ph/0606103.
- [14] M.-C. Chen and K. T. Mahanthappa, Phys. Lett. **B652**, 34 (2007), 0705.0714.
- [15] G. Altarelli, F. Feruglio, and C. Hagedorn, JHEP **03**, 052 (2008), 0802.0090.
- [16] S. F. King and M. Malinsky, Phys. Lett. **B645**, 351 (2007), hep-ph/0610250.
- [17] S.-L. Chen, M. Frigerio, and E. Ma, Nucl. Phys. **B724**, 423 (2005), hep-ph/0504181.
- [18] E. Ma, Mod. Phys. Lett. **A21**, 2931 (2006), hep-ph/0607190.
- [19] S. Morisi, M. Picariello, and E. Torrente-Lujan, Phys. Rev. **D75**, 075015 (2007), hep-ph/0702034.
- [20] W. Grimus and H. Kuhbock, Phys. Rev. **D77**, 055008 (2008), 0710.1585.
- [21] I. de Medeiros Varzielas and G. G. Ross, Nucl. Phys. **B733**, 31 (2006), hep-ph/0507176.
- [22] I. de Medeiros Varzielas, S. F. King, and G. G. Ross, Phys. Lett. **B648**, 201 (2007), hep-ph/0607045.
- [23] G. Altarelli and F. Feruglio, Nucl. Phys. **B720**, 64 (2005), hep-ph/0504165.
- [24] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. **B147**, 277 (1979).
- [25] I. de Medeiros Varzielas, S. F. King, and G. G. Ross, Phys. Lett. **B644**, 153 (2007), hep-ph/0512313.
- [26] E. Ma, Phys. Rev. **D70**, 031901 (2004), hep-ph/0404199.
- [27] K. S. Babu and X.-G. He (2005), hep-ph/0507217.
- [28] E. Ma, Mod. Phys. Lett. **A20**, 2601 (2005), hep-ph/0508099.

- [29] A. Zee, Phys. Lett. **B630**, 58 (2005), hep-ph/0508278.
- [30] G. Altarelli and F. Feruglio, Nucl. Phys. **B741**, 215 (2006), hep-ph/0512103.
- [31] X.-G. He, Y.-Y. Keum, and R. R. Volkas, JHEP **04**, 039 (2006), hep-ph/0601001.
- [32] G. Altarelli, F. Feruglio, and Y. Lin, Nucl. Phys. **B775**, 31 (2007), hep-ph/0610165.
- [33] E. Ma, Mod. Phys. Lett. **A22**, 101 (2007), hep-ph/0610342.
- [34] F. Bazzocchi, S. Kaneko, and S. Morisi, JHEP **03**, 063 (2008), 0707.3032.
- [35] L. Lavoura and H. Kuhbock, Eur. Phys. J. **C55**, 303 (2008), 0711.0670.
- [36] B. Brahmachari, S. Choubey, and M. Mitra, Phys. Rev. **D77**, 073008 (2008), 0801.3554.
- [37] M. Hirsch, S. Morisi, and J. W. F. Valle (2008), 0804.1521.
- [38] Y. Lin (2008), 0804.2867.
- [39] F. Feruglio, C. Hagedorn, Y. Lin, and L. Merlo (2008), 0807.3160.
- [40] F. Bazzocchi, S. Morisi, and M. Picariello, Phys. Lett. **B659**, 628 (2008), 0710.2928.
- [41] P. Minkowski, Phys. Lett. **B67**, 421 (1977).
- [42] M. Gell-Mann, P. Ramond, and R. Slansky, print-80-0576 (CERN), in Supergravity, P. van Nieuwenhuizen and D.Z. Freedman ed., North Holland, Amsterdam 1979, p.315.
- [43] T. Yanagida, Prog. Theor. Phys. **64**, 1103 (1980), in proc. of the *Workshop on the Unified Theory and Baryon Number in the Universe*, O.Sawada and A.Sugamoto eds., KEK report 79-18, 1979, p.95, Tsukuba, Japan.
- [44] S. L. Glashow (1979), in Cargese 1979, Proceedings, *Quarks and Leptons*, 687-713.
- [45] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [46] M. Magg and C. Wetterich, Phys. Lett. **B94**, 61 (1980).
- [47] J. Schechter and J. W. F. Valle, Phys. Rev. **D22**, 2227 (1980).
- [48] G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. **B181**, 287 (1981).
- [49] R. N. Mohapatra and G. Senjanovic, Phys. Rev. **D23**, 165 (1981).
- [50] G. Ross and M. Serna (2007), 0704.1248.
- [51] S. F. King and G. G. Ross, Phys. Lett. **B520**, 243 (2001), hep-ph/0108112.
- [52] E. Ma, Mod. Phys. Lett. **A17**, 627 (2002), hep-ph/0203238.
- [53] Z. G. Berezhiani, Phys. Lett. **B129**, 99 (1983).
- [54] S. Dimopoulos, Phys. Lett. **B129**, 417 (1983).
- [55] A. Davidson and K. C. Wali, Phys. Rev. Lett. **59**, 393 (1987).
- [56] Z. G. Berezhiani and R. Rattazzi, Phys. Lett. **B279**, 124 (1992).
- [57] L. Ferretti, S. F. King, and A. Romanino, JHEP **11**, 078 (2006), hep-ph/0609047.
- [58] Z. Berezhiani and F. Nesti, JHEP **03**, 041 (2006), hep-ph/0510011.
- [59] F. Feruglio, C. Hagedorn, Y. Lin, and L. Merlo (2007), hep-ph/0702194.
- [60] C. S. Lam, Phys. Lett. **B656**, 193 (2007), 0708.3665.
- [61] P. Nath and R. M. Syed, Phys. Lett. **B506**, 68 (2001), hep-ph/0103165.
- [62] P. Nath and R. M. Syed, Nucl. Phys. **B618**, 138 (2001), hep-th/0109116.
- [63] R. M. Syed (2005), hep-ph/0508153.
- [64] C. S. Aulakh and A. Girdhar, Int. J. Mod. Phys. **A20**, 865 (2005), hep-ph/0204097.
- [65] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac, and N. Okada, J. Math. Phys. **46**, 033505 (2005), hep-ph/0405300.